

This file has been cleaned of potential threats.

If you confirm that the file is coming from a trusted source, you can send the following SHA-256 hash value to your admin for the original file.

0e4004075efc9d6a0609579224b5c7513c97ea60aaaea150ef51c6e8bfe6d63

To view the reconstructed contents, please SCROLL DOWN to next page.

The text that follows is a PREPRINT.

Please cite as:

Fearnside, P.M. & Barbosa, R.I. 2004. Accelerating deforestation in Brazilian Amazonia:
Towards answering open questions. Environmental Conservation 31(1): 7-10.

ISSN: 0376-8929

Copyright: Cambridge University Press

The original publication is available at <http://journals.cambridge.org/> [<publisher link>](#)

Accelerating deforestation in Brazilian Amazonia: Towards answering open questions

PHILIP M. FEARNSIDE^{1*} AND REINALDO IMBROZIO BARBOSA¹
National Institute for Research in the Amazon (INPA), Av. André Araújo, 2936, C.P. 478,
69011-970 Manaus, Amazonas, Brazil

*Correspondence: Philip M. Fearnside, Tel: +55 92 643-1822 Fax: +55 92 642-8909 e-mail: pmfearn@inpa.gov.br

Running head: Deforestation in Amazonia

revised: 6 Oct. 2003

Total word count: 3442

In press: *Environmental Conservation*

SUMMARY

The annual rate of deforestation in Brazilian Amazonia jumped by an estimated 40% between 2001 and 2002. This increase is in addition to a 15% upward revision by the Brazilian government of the estimated rate for 2001. Examination of the data underlying these estimates and comparisons with other measurements indicates that important questions remain unanswered, especially in the state of Mato Grosso, where assessment of the effectiveness of a deforestation licensing and control program is critical to future efforts to contain forest destruction. The increase in deforestation rate in Mato Grosso in 2002 (23%) was less than half the increase in the remainder of Brazilian Amazonia (55%), indicating that the state government's deforestation control program may have had some effect.

Keywords: Deforestation, Amazonia, Brazil, Rainforest, Tropical forest, Remote sensing

INTRODUCTION

A series of open questions regarding Amazonian deforestation has become the focus of public attention following the June 2003 announcement of a 40% increase in the rate of Amazonian deforestation in 2002, as compared to the rate in 2001. This increase comes on top of an additional 15% increase in the estimated deforestation rate for 2001 (Fig. 1), which was announced at the same time (Brazil, INPE 2003). Here, we outline some of the doubts concerning the estimates and try to suggest ways that some of them might be resolved. The impetus of the new surge in forest clearing must be translated into both increased effort to quantify and understand the deforestation process and to take the sometimes politically and financially costly measures needed to contain it. The upsurge does not indicate that deforestation control measures have failed or that the deforestation process is inherently uncontrollable.

[Figure 1 here]

2001 DEFORESTATION

2001 ESTIMATES FOR THE LEGAL AMAZON

INPE's revision of the deforestation rate estimate for 2001 increased the amount from the 15,787 km²/year preliminary estimate released in 2002 to a final estimate of 18,165 km²/year (a 15% increase). Considerable fanfare had accompanied the announcement of the preliminary 2001 estimate as indicating a decrease from the rate in 2000, but the revised number indicates that no decrease took place (Fig. 1). The preliminary estimate for 2001 was based on 49 scenes where new clearing summed to 12,695 km² (Brazil, INPE 2001, page 20). A LANDSAT "scene" is a 185 km × 185 km area of the Earth's surface where the satellite makes an image each time it overflies the site. To produce the preliminary estimate, the 2001 deforestation total from the critical scenes was adjusted upward by 20% to represent the remainder of the 229 scenes covering Brazil's Legal Amazon region. This was based on the percentage differential in the previous year between the regional total from complete "wall-to-wall" coverage and the same sample of 49 "critical" scenes (Brazil, INPE 2002, page 20).

For some parts of the region the upward revision of the estimate for 2001 may still be an underestimate of the true clearing. For example, in the state of Roraima, an improbably low deforestation rate is indicated for the scene (231-60) covering part of counties of São Luiz do Anauá, São João da Baliza, Caroebe and Rorainópolis: clearing of only 49.7 km² by 2001. The real area is probably higher because these counties receive dozens of additional migrant families every month. Another example is the scene (233-58) covering the Trairão settlement area, which indicates zero deforestation in 2001. We visited this area in 1998, at which time the rapid advance of deforestation was readily apparent. Since that year, two new settlements have been established near Trairão. Apparently INPE was unable to obtain an image for this region (Brazil, INPE 2003). Of the 20 scenes needed to cover Roraima, 12 have no data for deforestation for the 2000-2001 period. Omissions of clearings should be leading to an underestimate of deforestation, rather than the overestimate that would be needed to explain the jump in INPE's deforestation estimate for the region.

2001 ESTIMATES FOR MATO GROSSO

Inconsistent results for 2001 in the state of Mato Grosso are a major concern. In Mato Grosso, FEMA (State Foundation for Environment), the state government's environmental agency, has been monitoring clearing at two-year intervals, with annual surveys being instituted in 2002 (Fearnside 2003). Since the FEMA monitoring program began, the FEMA and INPE estimates have never been in agreement, and have even differed in the direction of the discrepancy (Table 1).

[Table 1 here]

The term "deforestation" is used in a different way in Mato Grosso than in other contexts, such as the annual deforestation estimates released by INPE. INPE uses the term "*desflorestamento*" (a term invented by INPE) to refer to clearing of vegetation that corresponds to the forest and transition categories, and does not consider clearing of *cerrado* (savannah). In Mato Grosso state government usage, "deforestation" (*desmatamento*) refers to clearing of all three categories. The comparison between INPE and FEMA results in Table 1 excludes *cerrado* from the FEMA estimates.

INPE has suggested that the explanation of the differences between the INPE and FEMA results is due to differing definitions of the vegetation subject to deforestation: "forest" and "transition" (dos Santos 2002). However, considering the description of vegetation types defined by INPE as "forest" (Brazil, INPE 2001), the explanation of differing definitions is not particularly satisfactory. Only one relatively minor vegetation type ("*cerradão*," or high *cerrado*), designated "Sd" in the RADAMBRASIL map code (Brazil, Projeto RADAMBRASIL 1973-1983), is considered as forest by INPE and as *cerrado* (*i.e.*, non-forest) by FEMA. Measurements from RADAMBRASIL 1:250,000 and 1:1,000,000-scale maps indicate this vegetation type as totalling 26,083 km² in Mato Grosso, or only 3% of the state's vegetation (P.M. Fearnside & R.I. Barbosa 2003).

One possibility is that the prose descriptions of the vegetation types considered as "forest" by INPE do not match the areas considered as such in their geographical information system (GIS). INPE has not yet made its GIS coverages publicly available, although the intention to do so has been mentioned on various occasions over the years.

INPE's separation of the original (pre-Columbian) vegetation into "forest" (forest and transition) and non-forest (*cerrado*) is based on a mix of information types, including both RADAMBRASIL maps and appearance on LANDSAT imagery. Since INPE has never released estimates for the original forest areas by state derived using the definition of "forest" that INPE applies to the annual deforestation estimates (see reviews in Fearnside 1997, 2000), an estimate for Mato Grosso (Table 2) is derived from a coarse-scaled map (6 km²/pixel) released in 2003 (Brazil, MCT & MMA 2003). Measurements indicate that INPE considers "forest" (forest and transition) to be the original vegetation in 32.0% of the state's land area, substantially less than the 43.7% indicated by FEMA (Mato Grosso, FEMA 2002). This discrepancy, however, is in the wrong direction to explain the different results of the deforestation estimates: with less area considered as originally "forest" (*i.e.*, potentially subject to deforestation), INPE would be expected to find less, rather than more, deforestation than FEMA.

[Table 2 here]

In the case of FEMA, the limits between what was considered to have had original vegetation as forest, transition and *cerrado* were (possibly subject to change following a change of state government on 1 January 2003) defined exclusively on the basis of RADAMBRASIL maps. This was important as a legal matter in order to avoid opportunities for corruption influencing reclassification of land into categories where greater percentages of clearing would be legally permitted. Since 2000, federal requirements specify that the "legal reserve" that Brazil's 1965 "Forestry Code" requires in each property must cover 80% of the property in forest areas and 35% in *cerrado* areas. A Mato Grosso state government decision specifies 50% in the "transition" area. The FEMA definitions in use through 31 December 2002 are contained in digital maps that were distributed to forestry engineers who prepare applications for licensing (Mato Grosso, FEMA 2001a). Measurements from these maps indicate that 572,645 km², or 63.2% of the state, was originally either forest or transition by FEMA's definition (Mato Grosso, FEMA 2002). The difference between the FEMA and INPE definitions of forest+transition indicated by the maps is 14,804 km², but the size of the discrepancy is partially hidden by the 26,083 km² area of *cerradão* that is included as "forest" by INPE but not by FEMA. If *cerradão* is subtracted from the INPE forest area, the difference for the remaining "forest" vegetation is 40,887 km², an area the size of Belgium.

Another possible explanation could lie in differences in the dates of the images used. Neither the FEMA estimates nor the state-level estimates from INPE are normalized, for example by projecting the clearing for each scene to the end of the year. However, the images used are almost always from the most cloud-free period at the height of the dry season (after most felling is complete), and we believe it unlikely that different image dates could explain such a large discrepancy (up to 54.5%: Table 1).

2002 DEFORESTATION

2002 ESTIMATES FOR THE LEGAL AMAZON

The 25.5×10^3 km²/year preliminary figure for 2002 is not only higher than the rate for 2001, it is also higher than the 19×10^3 km²/year figure for 2002 that was leaked

to the press on 5 May 2003 (*Via Ecológica* 2003). Delays in releasing bad news have been a repeated pattern, as in the cases of the upturn of deforestation beginning in 1992 (not revealed until 1995), and the record peak of $29.1 \times 10^3 \text{ km}^2/\text{year}$ in 1995, not revealed until one month after the end of the December 1997 Kyoto conference on global warming (see Fearnside 1997, 2000). Release of the 2002 preliminary estimate was accompanied by the announcement that future deforestation estimates will be released without delays.

2002 ESTIMATES FOR MATO GROSSO

The location of changes in deforestation rate is important for understanding the increase in 2002. The deforestation in the 50 sample scenes used for preliminary estimate for deforestation in 2002 can be apportioned among the states, with any scenes that overlap more than one state being assigned to the state with the largest share of the scene (Table 3). Deforestation increased less in Mato Grosso (23.1%) than in the other states (54.5%). Particularly important are the larger increases in the two other major states for deforestation activity: Pará (35.4%) and Rondônia (50.5%). Tocantins, where deforestation rate decreased by 12.3%, is of limited importance for two reasons: Tocantins never had much forest because most of the state was originally *cerrado*, and, since much of the forest that did exist has already been cut, the decrease in deforestation rate is best explained by the dwindling of forest areas available for clearing. The smaller percentage increase in Mato Grosso relative to the increase in the total from sample scenes in other states suggests that the FEMA licensing and control program may have been having some effect on clearing. However, some of the lower rate of clearing in Mato Grosso is due to the diminishing areas of forest available for clearing. Mato Grosso is of particular importance because of the deforestation licensing and control program that was undertaken there by the state government's environmental agency over the 1999-2001 period (*e.g.*, Mato Grosso, FEMA 2001b).

[Tables 3 here]

CONCLUSIONS

Questions remain to be answered regarding deforestation estimates in Amazonia. Most important is clarification of the 2001 estimate for the state of Mato Grosso, where data from the state-government agency (FEMA) indicate that the deforestation rate decreased in forest and transition vegetation, while data from the federal agency (INPE) indicate that the rate increased. This clarification is needed to assess the effectiveness of Mato Grosso's program to license and control deforestation. Several indications point to an effect from the program, despite open questions regarding interpretation of the INPE data. For 2002, a major upsurge in deforestation occurred throughout Brazilian Amazonia, both inside and outside of Mato Grosso, but the increase was less in Mato Grosso than in the rest of the Amazonian region. Both INPE and FEMA should release their data and metadata (*i.e.*, documentation of the derivation of all intermediary results for each component of the estimates), allowing unrestricted public access via internet, so that the scientific community at large can examine the results, clarify outstanding discrepancies, and improve future monitoring procedures.

ACKNOWLEDGEMENTS

Our work is supported by the National Council for Scientific and Technological Development (CNPq)(Proc. 470765/01-1). We thank two reviewers for helpful comments.

References

- Brazil, INPE (Instituto Nacional de Pesquisas Espaciais). (2001) *Monitoramento da Floresta Amazônica Brasileira por Satélite/Monitoring of the Brazilian Amazon Forest by Satellite: 1999-2000*. São José dos Campos, São Paulo, Brazil: INPE.
- Brazil, INPE (Instituto Nacional de Pesquisas Espaciais). (2002) *Monitoramento da Floresta Amazônica Brasileira por Satélite/Monitoring of the Brazilian Amazon Forest by Satellite: 2000-2001*. São José dos Campos, São Paulo, Brazil: INPE.
- Brazil, INPE (Instituto Nacional de Pesquisas Espaciais). (2003) Taxas nos períodos 2000-2001 e 2001-2002 para 50 cenas críticas. (www document) URL http://www.obt.inpe.br/prodes/prodes_2002_2002.htm.
- Brazil, MCT (Ministério da Ciência e Tecnologia) & MMA (Ministério do Meio Ambiente). (2003) Mapeamento do Desmatamento da Amazônia com Imagens de Satélite. (www document) URL <http://www.obt.inpe.br/prodes.html>
- Brazil, Projeto RADAMBRASIL. (1973-1983) *Levantamento de Recursos Naturais*, Vols. 1-31. Rio de Janeiro, Brazil: Ministério das Minas e Energia, Departamento Nacional de Produção Mineral (DNPM).
- dos Santos, J.R. (2002) Visita técnica dos pesquisadores do INPE/S.J. Campos à FEMA-MT. Instituto Nacional de Pesquisas Espaciais (INPE), São José dos Campos, São Paulo, Brazil. Report to Ministério do Meio Ambiente (MMA), Secretaria de Coordenação da Amazônia (SCA), Sub-Programa de Recursos Naturais (SPRN), 31 July 2002. São José dos Campos, São Paulo, Brazil: INPE.
- Fearnside, P.M. (1997) Monitoring needs to transform Amazonian forest maintenance into a global warming mitigation option. *Mitigation and Adaptation Strategies for Global Change* **2**(2-3): 285-302.
- Fearnside, P.M. (2000) Effects of land use and forest management on the carbon cycle in the Brazilian Amazon. *Journal of Sustainable Forestry* **12**(1/2): 79-97.
- Fearnside, P.M. (2003) Deforestation control in Mato Grosso: A new model for slowing the loss of Brazil's Amazon forest. *Ambio* **32**(5): 343-345.
- Fearnside, P.M. & Barbosa, R.I. (2003) Avoided deforestation in Amazonia as a global warming mitigation measure: The case of Mato Grosso. *World Resource Review* (in press).
- Mato Grosso, Fundação Estadual do Meio Ambiente (FEMA). (2001a) Sistema de licenciamento ambiental único, Versão 1.0. (CD-ROM) Cuiabá, Mato Grosso, Brazil: FEMA, Diretoria de Recursos Florestais (DRF).

Mato Grosso, Fundação Estadual do Meio Ambiente (FEMA). (2001b) Environmental Control System on Rural Properties in Mato Grosso. Cuiabá, Mato Grosso, Brazil: FEMA.

Mato Grosso, Fundação Estadual do Meio Ambiente (FEMA). (2002) Dinâmica do desmatamento no Estado de Mato Grosso 2001. Cuiabá, Mato Grosso, Brazil: FEMA.

Via Ecológica. (2003) “Satélites mostram que agricultura ajudou a destruir mais floresta na Amazônia.” *Via Ecológica* 5 May 2003 (www document) URL <http://www.viaecologica.com.br>

FIGURE LEGEND

Figure 1 –Estimates of deforestation in Brazil's Legal Amazon region (Brazil, INPE 2002, 2003). The 2002 estimate is preliminary.

Table 1 Comparison of FEMA and INPE deforestation data for Mato Grosso for forest and transition (forest-savannah ecotone) only; *cerrado* clearing, which is not monitored by INPE, is excluded from the FEMA values for this comparison.

Two-year period*	INPE (ha/ year)	FEMA (ha/year)	Discrepancy (ha/year)	(INPE-FEMA) (%)
1996-97	590,700	770,130	-179,430	-30.4
1998-99	671,450	638,066	+33,384	+5.0
2000-01	911,900	414,507	+497,393	+54.5

*Periods are standardized to the two-year “biennium” that FEMA estimates adopted through 2001. FEMA and INPE use the term ‘biennium’ (*biênio*) differently: FEMA to refer to a 24-month period and INPE to refer to a 12-month period (dry season to dry season) spanning two calendar years.

Table 2 Original vegetation and clearing in Mato Grosso

Agency	Original area of vegetation		Area cleared by 2001	
	(km ²)	(% of total vegetation)	(km ²)	(% of original area)
<i>Forest + transition^a</i>				
INPE ^b	509,146	59.5	163,005	32.0
FEMA	523,950	61.2	228,986	43.7
Discrepancy	-14,804	-1.7	-65,982	-11.7
(INPE-FEMA)	(-2.9%)		(-40.5%)	
<i>Cerrado^a</i>				
INPE ^b	347,227	40.5	--- ^c	--- ^c
FEMA	332,423	38.8	124,121	37.3
Discrepancy	+14,804	+1.7		
(INPE-FEMA)	(+4.3%)			

- (a) INPE includes *cerradão* (Sd), estimated at 26,083 km² (see text), in forest + transition; FEMA includes *cerradão* in the *cerrado* category.
- (b) INPE areas calculated from percentages multiplied by FEMA area for total vegetation. INPE percentages are measured from a map released in 2003 (Brazil, MCT and MMA 2003) ignoring 1.4% of area not classified. The forest + transition cleared area by 2001 derived here from the INPE map is greater than the 151.633 km² implied by past cumulative clearing (Brazil, INPE 2002) and the rate value released in 2003 (Brazil, INPE 2003).
- (c) INPE does not measure *cerrado* clearing.

Table 3 Deforestation in 50 sample scenes. Data from Brazil, INPE (2003).

State	Number of scenes	Year		(%)
		2001	2002	Change
Mato Grosso	19	6,250	7,695	+23.1
Rondônia	9	2,628	3,956	+50.5
Pará	13	3,388	4,589	+35.4
Maranhão	4	751	1,561	+108.0
Amazonas	2	171	376	+119.6*
Tocantins	2	454	398	-12.3
Acre	1	106	704	+566.8
Total	50	13,747	19,279	+40.2

* In one of the scenes assigned to Amazonas, most of this increase is probably in the portion of the scene located in Rondônia.

Figure 1

