

This file has been cleaned of potential threats.

If you confirm that the file is coming from a trusted source, you can send the following SHA-256 hash value to your admin for the original file.

f95082e36bf998b6f964418cbb1b24224947edb0317158e53a2b30a554f7b90c

To view the reconstructed contents, please SCROLL DOWN to next page.

The text that follows is a PREPRINT

O texto que segue é um PREPRINT.

Please cite as:

Favor citar como:

**Laurance, W.F.; J.L.C. Camargo, P.M.
Fearnside, T.E. Lovejoy, G.B.
Williamson, R.C.G. Mesquita, C.F.J.
Meyer, P.E.D. Bobrowiec, & S.G.W.
Laurance. 2016. An Amazonian forest
and its fragments as a laboratory of
global change. pp. 407-440. In: L. Nagy,
B. Forsberg, P. Artaxo (eds.) *Interactions
Between Biosphere, Atmosphere and
Human Land Use in the Amazon Basin..*
Springer (Ecological Studies 227), Berlin,
Alemanha. 478 pp.**

doi: 10.1007/978-3-662-49902-3_16

ISBN (Hard Cover): 978-3-662-49902-3;

ISBN (Ebook): 978-3-662-49900-9.

Copyright: Springer Science + Business Media

The original publication is available from:

A publicação original está disponível de:

<http://www.springer.com/us/book/9783662499009>

Springer
Tiergartenstraße 17
69121 Heidelberg
Germany

An Amazonian forest and its fragments as a laboratory of global change

William F. Laurance^{1,*}, José L. C. Camargo², Philip M. Fearnside³, Thomas E. Lovejoy^{2,4},
G. Bruce Williamson⁵, Rita C. G. Mesquita⁶, Christoph F. J. Meyer^{2,7}, Paulo E. D.
Bobrowiec^{2,8} & Susan G. W. Laurance^{1,*}

¹Centre for Tropical Environmental and Sustainability Science (TESS) and College of Marine and Environmental Science, James Cook University, Cairns, Queensland 4878, Australia,

*corresponding authors: bill.laurance@jcu.edu.au and susan.laurance@jcu.edu.au

²Biological Dynamics of Forest Fragments Project, National Institute for Amazonian Research (INPA) and Smithsonian Tropical Research Institute, Manaus, AM 69067-375, Brazil, zeluiscamargo@gmail.com

³Department of Environmental Dynamics, National Institute for Amazonian Research (INPA), Manaus, AM 69067-375, Brazil, phillip.fearnside@gmail.com

⁴Department of Environmental Science and Policy, George Mason University, Fairfax, VA 22030, USA, tlovejoy@unfoundation.org

⁵Department of Biological Sciences, Louisiana State University, Baton Rouge, LA, 70803, USA, btwill@lsu.edu

⁶Long-term Ecological Research (PELD), National Institute for Amazonian Research (INPA), Manaus, AM 69067-375, Brazil, rita@buriti.com.br

⁷Center for Environmental Biology, University of Lisbon, Campo Grande C2, 1749-016 Lisbon, Portugal; E-mail: cmeyer@fc.ul.pt

⁸Coordination of Biodiversity, National Institute for Amazonian Research (INPA), Manaus, AM 69067-375, Brazil, paulobobro@gmail.com

Abstract

We synthesize findings from the world's largest and longest-running experimental study of habitat fragmentation, in central Amazonia. Over the past 36 years, 11 forest fragments ranging from 1-100 ha in size have experienced a wide array of ecological changes. Edge effects have been a dominant driver of fragment dynamics, strongly affecting forest microclimate, tree mortality, carbon storage, and fauna. The matrix of vegetation surrounding fragments has changed markedly over time (evolving from large cattle pastures to mosaics of abandoned pasture and secondary regrowth forest), and this, in turn, has strongly influenced the dynamics of fragments and faunal communities. Both rare weather events and apparent global-change drivers have significantly influenced forest structure and dynamics across the entire study area, both in forest fragments and in nearby intact forest. Such large-scale drivers are likely to interact synergistically with habitat fragmentation.

Keywords: Amazonia, Biodiversity, Climate change, Edge effects, Ecosystem services, Environmental synergisms, Global change, Habitat fragmentation, Rainforest

1. Introduction

The Biological Dynamics of Forest Fragments Project (BDFFP) is the world's largest and longest-running experimental study of habitat fragmentation¹ (Box 1). Located in central Amazonia near the city of Manaus, the BDFFP has evolved since its inception in 1979 into a major epicenter for long-term research. The BDFFP's research mission has gradually broadened to include not only long-term studies of forest fragmentation but also important work on global-change phenomena and a variety of basic research topics.

Here we highlight some key contributions of this singular project to the study of land-use change and regional- and global-scale drivers in central Amazonia, at the heart of the world's largest tropical forest.

1.1 Amazonia and global change

Amazonia stands at the intersection of several key questions for global change, both for study and for action. It is believed to be one of the regions that will be most impacted by projected climate changes (Salazar et al. 2007, Dai 2012, IPCC 2013). It has the potential to contribute significantly to efforts to mitigate climate change during the narrow window of time that we have to avert 'dangerous' global warming (Fearnside 2000, 2012). It is also one of the places where sharply reducing greenhouse-gas emissions—by limiting forest loss and degradation—could deliver the greatest global benefits for humankind (Stickler et al. 2009).

The rapid loss and fragmentation of old-growth forests are among the greatest threats to tropical biodiversity (Lovejoy et al. 1986, Sodhi et al. 2004, Laurance and Peres 2006, Gibson et al. 2011). More than half of all remaining tropical forest occurs in the Amazon Basin, which is being seriously altered by large-scale agriculture (Fearnside, 2001a, Gibbs et al. 2010), industrial logging (Asner et al. 2005), proliferating roads and energy infrastructure (Laurance et al. 2001a, Fearnside 2002, 2007, Killeen 2007), increasing biofuel production (Butler and Laurance 2009), and oil, gas, and mining developments (Finer et al. 2008).

The exploitation of Amazonia is driving forest fragmentation on a vast spatial scale. By the early 1990s, the area of Amazonian forest that was fragmented ($<100 \text{ km}^2$) or vulnerable to edge effects ($<1 \text{ km}$ from edge) was over 150% greater than the area that had been deforested (Skole and Tucker 1993). From 1999 to 2002, deforestation and logging in Brazilian Amazonia respectively created $\sim 32,000$ and $\sim 38,000 \text{ km}$ of new forest edge annually (Broadbent et al. 2008). Prevailing land uses in Amazonia, such as cattle ranching and small-scale farming, produce landscapes dominated by small ($<400 \text{ ha}$) and irregularly shaped forest fragments (Cochrane and Laurance 2002, Broadbent et al. 2008). Such fragments are highly vulnerable to edge effects, fires, and other deleterious consequences of forest fragmentation (Laurance et al. 2002, Barlow et al. 2006, Cochrane and Laurance 2008).

While model predictions for future climate in Amazonia vary considerably, it is expected that parts of the region will be hotter and drier under expected global warming (Dai 2012, IPCC 2013). What this warming portends for affected areas of Amazonian forest is a matter of some controversy. Disastrous die-offs projected by the UK Meteorological Office Hadley Centre at atmospheric CO_2 concentrations about twice those in the pre-industrial atmosphere (Cox et al. 2000, 2004) have now been countered by a new model version from the same group indicating the Amazon forest almost entirely intact, even with up to four times the pre-industrial CO_2 concentration (Cox et al. 2013, Good et al. 2013, Huntingford et al. 2013). The main difference is inclusion of CO_2 -fertilization effects, making the trees grow faster, resist stress better, and close their stomata more frequently such that they use and need less water.

1.2 Contributions of the BDFFP to global-change research

The BDFFP, with 36 years of research in fragmented and continuous forest in central Amazonia (Box 1; Fig. 1), has been contributing to quantifying the interactions of land use and global climate change. BDFFP studies have assessed the vulnerability of the forest to changes in meteorological parameters (Laurance et al. 2009a), including those that are aggravated by fragmentation (Laurance 2004). The long-term monitoring of thousands of individual forest trees, and of populations of various other plant and animal species in the same locations, provides the potential for early detection of global environmental changes.

The BDFFP is a source of invaluable long-term datasets. These include high-quality estimates of Amazon forest biomass and carbon stocks (Phillips et al. 1998, Baker et al. 2004). The project also contributes greatly to knowledge of the diversity of species and their relationships in an Amazon forest ecosystem (Laurance et al. 2010a, ter Steege et al. 2013). Biodiversity and ecosystem processes represent part of what is lost when the forest is destroyed or degraded, whether by direct human action, by climate change, or by the interaction of both together. Understanding these processes is essential for assessing not only the vulnerability of forests, but also their potential resilience in the face of global change and the rate of recovery following perturbation (Williamson et al. 2014).

The BDFFP has made a substantial contribution to debate over climatic influences on the Amazon via its monitoring of lianas and forest dynamics (Laurance et al. 2014a, 2014b). Lianas evidently make better use of rising CO₂ than do trees (e.g., Condon et al. 1992), and contribute significantly to tree damage and mortality (Ingwell et al. 2010). They also form heavily vine-dominated ‘liana forests’ in drier parts of Amazonia (Fearnside 2013). BDFFP plots show a marked increase in liana abundance and biomass between censuses in 1997-99 and 2012 (Laurance et al. 2014). Liana increases have also been found in tropical forests in western Amazonia, central America, the Guianas, and elsewhere, with rising CO₂ levels being one of the more likely explanations (see Laurance et al. 2014 and references therein). This negative effect of CO₂ enrichment is not included in the Hadley Centre models, and would likely cancel out some of the benefits indicated in a high-CO₂ future.

BDFFP data have helped to identify the direct effects of a warmer, drier climate on the forest. The microclimate at forest edges is significantly hotter and drier than that in the continuous forest (Kapos 1989, Kapos et al. 1993, Camargo and Kapos 1995). Canopy trees are vulnerable to changing microclimates on forest edges during the dry season, with desiccation detected up to 2 km from clearings (Briant et al. 2010). At the BDFFP, edge-associated tree mortality and ‘biomass collapse’ have been extensively documented (Laurance et al. 1997, 1998, 2000, Nascimento and Laurance 2004). Because the entire forest can be expected to face comparable conditions under projected climate change, the dead trees in the BDFFP fragment edges stand as a clear warning of the power of these changes.

Better estimates of how the forest will fare under changed climate are essential for many reasons, including providing the scientific basis needed to convince both world leaders and the general public that containing climate change is worth the cost. But just as basic is the question: what should we do about climate change once the world finally decides to act? The role of tropical forests is critical to this debate, as they contain a large stock of carbon that could either be substantially released by deforestation, logging, and fire, or conserved for their crucial environmental values. The ways that avoiding these emissions could be incorporated into global mitigation efforts, how carbon benefits would be rewarded, and how they should be calculated have been the subject of longstanding controversy dividing environmental groups, national governments, and scientists (see Fearnside 2001b, 2012).

One aspect of this discussion to which the BDFFP makes an important contribution is in reducing the uncertainty surrounding biomass and carbon-stock estimates for Amazon forest. The BDFFP tree survey is much more complete than, for example, the 3000 1-ha plots surveyed by the RADAMBRASIL Project (Nogueira et al. 2008). The BDFFP has much

better species identifications and includes data on other forest components, such as palms, lianas, strangler figs, understory vegetation, and dead vegetation (necromass). Correct species identification allows better matching with plant functional traits such as wood density and tree form (e.g., Fearnside 1997, Nogueira et al. 2005, 2007, Chave et al. 2006).

Crucially, the BDFFP forest data allow one to see the variability in biomass from one hectare to another. The mean aboveground biomass of live trees across 69 1-ha plots was $356 \pm 47 \text{ Mg ha}^{-1}$ (Laurance et al. 1999). This great variability indicates the need for many plots, rather than relying on only a few plots of 1 ha or less scattered around the region as the basis for calibrating satellite imagery for biomass mapping and for estimating greenhouse-gas emissions from deforestation (see Fearnside 2014).

2. Long-term studies of forest fragmentation

The BDFFP's original mission focuses on assessing the effects of forest fragmentation on Amazonian forests and fauna, and on important ecological and ecosystem processes. Here we summarize some key conservation lessons that have been gleaned to date.

2.1. Sample effects are important in Amazonia

Many species in Amazonian forests are rare or patchily distributed. This phenomenon is especially pronounced in the large expanses of the basin that overlay heavily weathered, nutrient-poor soils (e.g. Radtke et al. 2008), where resources such as fruits, flowers, and nectar are scarce and plants are heavily defended against herbivore attack (Laurance 2001). Herein lies a key implication for understanding forest fragmentation: given their rarity, many species may be absent from fragments not because their populations have vanished, but because they were simply not present at the time of fragment creation—a phenomenon termed the 'sample effect' (Wilcox and Murphy 1985). Such sample effects are the hypothesized explanation for the absence of many rare understory bird species from fragments (Ferraz et al. 2007). In addition, many beetles (Didham et al. 1998a), bats (Sampaio et al. 2003), ant-defended plants (Bruna et al. 2005), and trees (Bohlman et al. 2008, Laurance et al. 2010b) at the BDFFP exhibit high levels of habitat specialization or patchiness. In a region where rarity and patchy distributions of species are the norm, sample effects appear to play a major role in structuring fragmented communities. Given these sample effects, nature reserves will have to be especially large to sustain viable populations of rare species (Lovejoy and Oren 1981, Laurance 2005, Peres 2005, Radtke et al. 2008).

2.2. Fragment size is vital

Although fragments range from just 1–100 ha in the BDFFP study area, understanding fragment-area effects has long been a central goal of the project (Lovejoy and Oren 1981, Lovejoy et al. 1984, 1986). The species richness of many organisms declines with fragment area, even with constant sampling effort across all fragments. Such declines are evident in leaf bryophytes (Zartman 2003), tree seedlings (Benítez-Malvido and Martinez-Ramos 2003a), palms (Scariot 1999), understory insectivorous birds (Stratford and Stouffer 1999; Ferraz et al. 2007), gleaning animal-eating bats (Sampaio 2000, Meyer et al. unpublished data), primates (Gilbert and Setz 2001, Boyle and Smith 2010a), and larger herbivorous mammals (Timo 2003), among others. For these groups, smaller fragments are often unable to support viable populations and deleterious edge effects—ecological changes associated with the abrupt, artificial edges of forest fragments—can also rise sharply in intensity (Didham et al. 1998a). A few groups, such as ant-defended plants and their ant mutualists, show no significant decline in diversity with fragment area (Bruna et al. 2005).

Fragment size also influences the rate of species losses, with smaller fragments losing species more quickly (Lovejoy et al. 1986, Stouffer et al. 2008). Assuming the surrounding

matrix is hostile to bird movements and precludes colonization, Ferraz et al. (2003) estimated that a 1000-fold increase in fragment area would be needed to slow the rate of local species extinctions by 10-fold. Even a fragment of 10,000 ha in area would be expected to lose a substantial part of its bird fauna within one century (Ferraz et al. 2003). Similarly, mark-recapture data suggest that very large fragments will be needed to maintain fully intact assemblages of some faunal groups, such as ant-following birds, which forage over large areas of forest (Van Houtan et al. 2007).

3. Edge effects

3.1. Forest hydrology is disrupted

The hydrological regimes of fragmented landscapes differ markedly from those of intact forest (Kapos 1989). Pastures or crops surrounding fragments have much lower rates of evapotranspiration than do forests because they have far lower leaf area and thus less rooting depth. Additionally, such clearings are hotter and drier than forests (Camargo & Kapos 1995). Field observations and heat-flux simulations suggest that desiccating conditions can penetrate up to 100–200 m into fragments from adjoining clearings (Malcolm 1998; Didham and Lawton 1999). Further, streams in fragmented landscapes experience greater temporal variation in flows than do those in forests, because clearings surrounding fragments have less evapotranspiration and rainfall interception by vegetation (Trancoso 2008). Free runoff promotes localized flooding in the wet season and stream failure in the dry season, with potentially important impacts on aquatic invertebrates (Nessimian et al. 2008) and other organisms.

Forest fragmentation also can alter low-level atmospheric circulation, which in turn affects local cloudiness and rainfall. The warm, dry air over clearings tends to rise, creating zones of low air pressure. The relatively cool, moist air over forests is drawn into this vacuum (Avisar and Schmidt, 1998). As it warms it also rises and forms convectional clouds over the clearing, which can lead to localized thunderstorms (Avisar and Liu 1996). In this way, clearings of a few hundred hectares or more can draw moisture away from nearby forests (Laurance 2004a, Cochrane and Laurance 2008). In Eastern Amazonia, satellite observations of canopy-water content suggest such desiccating effects typically penetrate 1.0–2.7 km into fragmented forests (Briant et al. 2010). This moisture-robbing function of clearings, in concert with frequent burning in adjoining pastures, could help explain why fragmented forests are so vulnerable to destructive, edge-related fires (Cochrane and Laurance 2002, 2008).

3.2. Edge effects often dominate fragment dynamics

Edge effects are among the most important drivers of ecological change in the BDFFP fragments. The distance to which different edge effects penetrate into fragments varies widely, ranging from <10 to 300 m at the BDFFP (Laurance et al. 2002) and considerably further (at least 2–3 km) in areas of the Amazon where edge-related fires are common (Cochrane and Laurance 2002, 2008; Briant et al. 2010).

Edge phenomena are remarkably diverse (Fig. 2). They include increased desiccation stress, wind shear, and wind turbulence that sharply elevate rates of tree mortality and damage (Laurance et al. 1997, 1998a). These in turn cause wide-ranging alterations in the community composition of trees (Laurance et al. 2000, 2006a, 2006b) and lianas (Laurance et al., 2001b). Such stresses may also reduce germination (Bruna 1999) and establishment (Uriarte et al., 2010) of shade-tolerant plant species in fragments, leading to dramatic changes in the composition and abundance of tree seedlings (Benítez-Malvido 1998, Benítez-Malvido and Martinez-Ramos 2003a).

Many animal groups, such as numerous bees, wasps, flies (Fowler et al. 1993), beetles

(Didham et al. 1998a, 1998b), ants (Carvalho and Vasconcelos 1999), butterflies (Brown and Hutchings 1997), understory birds (Quintela 1985, Laurance 2004b), and gleaning animal-eating bats (Rocha et al. 2013) decline in abundance near fragment edges. Negative edge effects are apparent even along forest roads (20–30 m width) in large forest tracts. Among understory birds, for example, five of eight foraging guilds declined significantly in abundance within 70 m of roads, whereas tree mortality increased and canopy cover declined (Laurance 2004b).

Some groups of organisms remain stable or even increase in abundance near edges. Leaf bryophytes (Zartman and Nascimento 2006), wandering spiders (*Ctenus* spp; Rego et al. 2007, Mestre and Gasnier 2008), and many frogs (Gascon 1993) show no significant response to edges. Organisms that favor forest ecotones or disturbances, such as many species of gap-favoring and frugivorous birds (Laurance 2004b), hummingbirds (Stouffer and Bierregaard 1995a), frugivorous bats that exploit early successional plants (Sampaio 2000), light-loving butterflies (Leidner et al. 2010), and fast-growing lianas (Laurance et al. 2001b), increase in abundance near edges, sometimes dramatically.

3.3. Edge effects are cumulative

BDFFP research provides strong support for the idea that two or more nearby edges create more severe edge effects than does just one (Fig. 3). This conclusion is supported by studies of edge-related changes in forest microclimate (Kapos 1989, Malcolm 1998), vegetation structure (Malcolm 1994), tree mortality (Laurance et al. 2006a), abundance and species richness of tree seedlings (Benítez-Malvido 1998, Benítez-Malvido and Martinez-Ramos 2003a), liana abundance (Laurance et al. 2001b), and the density and diversity of disturbance-loving pioneer trees (Laurance et al. 2006a, 2006b, 2007). The additive effects of nearby edges could help to explain why small (<10 ha) or irregularly shaped forest remnants are often so severely altered by forest fragmentation (Zartman 2003, Laurance et al. 2006a).

3.4. Edge age, structure, and adjoining vegetation influence edge effects

When a forest edge is newly created it is open to fluxes of wind, heat, and light, creating sharp edge-interior gradients in forest microclimate that stress or kill many rainforest trees (Lovejoy et al. 1986, Sizer and Tanner 1999). As the edge ages, however, proliferating vines and lateral branch growth tend to ‘seal’ the edge, making it less permeable to microclimatic changes (Camargo and Kapos 1995, Didham and Lawton 1999). Tree death from microclimatic stress is likely to decline over the first few years after edge creation (D’Angelo et al. 2004) because the edge becomes less permeable, because many drought-sensitive individuals die immediately, and because surviving trees may acclimate to drier, hotter conditions near the edge (Laurance et al. 2006a). Tree mortality from wind turbulence, however, probably increases as the edge ages and becomes more closed because, as suggested by wind-tunnel models, downwind turbulence increases when edges are less permeable (Laurance 2004a).

Regrowth forest adjoining fragment edges can also lessen edge-effect intensity. Microclimatic alterations (Didham and Lawton 1999), tree mortality (Mesquita et al. 1999), and edge avoidance by understory birds (Develey and Stouffer 2001, Laurance 2004b, Laurance et al. 2004), and gleaning bats that feed on invertebrates and small vertebrates (Meyer et al., 2013) are all reduced substantially when forest edges are buffered by adjoining regrowth forest, relative to edges adjoined by cattle pastures.

4. Isolation and matrix effects

4.1. Matrix structure and composition affect fragments

Secondary forests have gradually overtaken most pastures in the BDFFP landscape. This re-

growth lessens the effects of fragmentation for some taxa as the matrix becomes less hostile to faunal use and movements. Several species of insectivorous birds that had formerly disappeared have recolonized fragments as the surrounding secondary forest grew back (Stouffer and Bierregaard 1995b). The rate of local extinctions of birds has also declined (Stouffer et al. 2008). Similarly, gleaning animal-eating bats, which occurred at low abundances in fragments (Sampaio 2000) and in secondary regrowth (Bobrowiec and Gribel 2010) 10-15 years ago, have since increased in response to matrix regeneration (Meyer et al. 2013). A number of other species, including certain forest spiders (Mestre and Gasnier 2008), dung beetles (Quintero and Roslin 2005), euglossine bees (Becker et al. 1991), and monkeys such as red howlers, bearded sakis, and brown capuchins (Boyle and Smith 2010a) have recolonized some fragments.

The surrounding matrix also has a strong effect on plant communities in fragments by mediating certain edge effects (see above), influencing the movements of pollinators (Dick 2001, Dick et al. 2003) and seed dispersers (Jorge 2008, Bobrowiec and Gribel 2010, Boyle and Smith 2010a), and strongly affecting the seed rain that arrives in fragments. For instance, pioneer trees regenerating in fragments differed strikingly in composition between fragments surrounded by *Cecropia*-dominated regrowth and those encircled by *Vismia*-dominated regrowth (Nascimento et al. 2006). In this way plant and animal communities in fragments could come to mirror to some extent the composition of the surrounding matrix (Laurance et al. 2006a, 2006b), a phenomenon observed elsewhere in the tropics (e.g. Janzen 1983, Diamond et al. 1987).

4.2 Matrix is affected by history and forest proximity

Land-use history is a primary driver of secondary succession in the Central Amazon, resulting in the establishment of distinct trajectories differing in structure, composition, biomass, and dynamics (Mesquita et al. 1999, Williamson et al. 2014). Intensive use with prescribed fire to maintain pastures compromises the regenerative potential of land which, once abandoned, is colonized by few species and dominated by the genus *Vismia*, resulting in secondary forests that are depauperate in richness and stalled in succession. Where land use has been less intensive, a more diverse vegetation, dominated by the genus *Cecropia* colonizes, fostering relatively rapid plant succession.

Plant density and species diversity in secondary forests decrease with distance from forest edge, and are significantly different between *Vismia* and *Cecropia* dominated secondary forests. These differences were initially attributed to differential seed dispersal limitations (Mesquita et al. 2001, Puerta, 2002). Wieland et al. (2011), however, showed that the seed rain was similar for both types of second-growth and dominated by pioneer species, with only the occasional presence of mature forest species, even very close to forest edges. These results point to other relevant processes affecting plant establishment, such as seed consumption, germination success, and seedling herbivory (Wieland et al. 2011, Massoca et al. 2013).

4.3 Even narrow clearings are harmful

Many Amazonian species avoid clearings, and even a forest road can be an insurmountable barrier for some. A number of understory insectivorous birds exhibit depressed abundances near forest roads (20–40 m width) (Laurance 2004b) and strongly inhibited movements across those roads (Laurance et al. 2004). Experimental translocations of resident adult birds reveal such bird species will cross a highway (50–75 m width) but not a small pasture (250 m width) to return to their territory (Laurance and Gomez 2005). Individuals of other vulnerable species, however, have traversed clearings to escape from small fragments to larger forest areas (Harper 1989, Van Houtan et al. 2007). Captures of understory birds declined

dramatically in fragments when a 100 m-wide swath of regrowth forest was cleared around them, suggesting that species willing to traverse regrowth would not cross clearings (Stouffer et al. 2006).

Aside from birds, clearings of just 100–200 m width can evidently reduce or halt the movements of many forest-dependent organisms (Laurance et al. 2009b), ranging from herbivorous insects (Fáveri et al. 2008), euglossine bees (Powell and Powell 1987), and dung beetles (Klein 1989) to the spores of epiphyllous lichens (Zartman and Nascimento 2006, Zartman and Shaw 2006). Narrow clearings can also provide invasion corridors into forests for exotic and nonforest species (Gascon et al. 1999; Laurance et al. 2009b).

5. Landscape dynamics

5.1. Rare disturbances can leave lasting legacies

Rare events such as windstorms and droughts have strongly influenced the ecology of fragments. Rates of tree mortality rose abruptly in fragmented (Laurance et al., 2001c) and intact forests (Williamson et al. 2000, Laurance et al. 2009a) in the year after the intense 1997 El Niño drought. Such pulses of tree death help drive changes in the floristic composition and carbon storage of fragments (Laurance et al. 2007). Leaf-shedding by drought-stressed trees also increases markedly during droughts, especially within ~60 m of forest edges (Laurance and Williamson 2001). The additional litter increases the susceptibility of fragments to intrusion by surface fires (Cochrane and Laurance 2002, 2008).

Intense windblasts from convectional thunderstorms have occasionally strafed parts of the BDFFP landscape and caused intense forest damage and tree mortality, especially in the fragments. Fragments in the easternmost cattle ranch at the BDFFP have had substantially lower rates of tree mortality than did those in the other two ranches, because the former have so far escaped windstorms (Laurance et al. 2007). These differences have strongly influenced the rate and trajectory of change in tree-community composition in fragments (Laurance et al. 2006b). Hence, by altering forest dynamics, composition, structure, and carbon storage, rare disturbances have left an enduring imprint on the ecology of fragmented forests.

5.2. Fragments are hyperdynamic

The BDFFP fragments experience exceptionally large variability in population and community dynamics, relative to intact forest, despite being largely protected from ancillary human threats such as fires, logging, and overhunting. Being a small resource base, a habitat fragment is inherently vulnerable to stochastic effects and external vicissitudes. Species abundances can fluctuate dramatically in small communities, especially when immigration is low and disturbances are frequent (Hubbell 2001). Edge effects, reduced dispersal, external disturbances, and changing herbivore or predation pressure can all elevate the dynamics of plant and animal populations in fragments (Laurance 2002, 2008).

Many examples of hyperdynamism have been observed in the BDFFP fragments. Some butterfly species have experienced dramatic population irruptions in response to a proliferation of their favored host plants along fragment margins (Brown and Hutchings 1997), and butterfly communities in general are hyperdynamic in fragments (Leidner et al. 2010). Bat assemblages also show pronounced species turnover, particularly in 1-ha fragments (Meyer et al. 2013). Streamflows are far more variable in fragmented than forested watersheds (Trancoso 2008). Rates of tree mortality and recruitment are chronically elevated in fragments (Laurance et al. 1998a, b), with major pulses associated with rare disturbances (see above). Further, tree species disappear and turn over far more rapidly in fragments than intact forest, especially within ~100 m of forest margins (Laurance et al. 2006b). These and many other instabilities plague small, dwindling populations in the BDFFP fragments.

5.3. Fragments in different landscapes diverge

An important insight is that different fragmented landscapes— even those as alike as the three large cattle ranches in the BDFFP, which have very similar forests, soils, climate, fragment ages, and land-use histories—can diverge to a surprising degree in species composition and dynamics. Although spanning just a few dozen kilometers, the three ranches are following unexpectedly different trajectories of change.

At the outset, small initial differences among the ranches multiplied into much bigger differences. Parts of the western and eastern ranches were cleared in 1983, when an early wet season prevented burning of the felled forest. Tall, floristically diverse *Cecropia*-dominated regrowth quickly developed in these areas, whereas areas cleared in the years just before or after became cattle pastures or, eventually, scrubby *Vismia*-dominated regrowth (Williamson and Mesquita 2001). These different successional trajectories manifested, for instance, as distinct differences in bat assemblages, whereby *Cecropia*-dominated regrowth retained a considerable fraction of forest-specialist bat species found in continuous forest compared to *Vismia* regrowth (Bobrowiec and Gribel 2010). As discussed above, the differing matrix vegetation strongly affected the dynamics of plant and animal communities in the nearby fragments. These differences were magnified by subsequent windstorms, which heavily damaged most fragments in the central and western ranches, yet left fragments in the eastern ranch unscathed. Even identically sized fragments in the three ranches have had remarkably different dynamics and vectors of compositional change (Laurance et al. 2007).

The apparently acute sensitivity of fragments to local landscape and weather dynamics—even within a study area as initially homogeneous as ours—prompted us to propose a ‘landscape-divergence hypothesis’ (Laurance et al. 2007). We argue that fragments within the same landscape tend to have similar dynamics and trajectories of change in species composition, which will often differ from those in other landscapes. Over time, this process will tend to homogenize fragments in the same landscape, and promote ecological divergence among fragments in different landscapes. Evidence for this hypothesis is provided by tree communities in our fragments, which appear to be diverging in composition among the three cattle ranches (Fig. 4). Pioneer and weedy trees are increasing in all fragments, but the composition of these generalist plants and their rate of increase differ markedly among the three ranches (Scariot 2001, Laurance et al. 2006a, 2007, Nascimento et al. 2006). This same pattern of landscape homogenization within ranches can also be seen for bat assemblages in the secondary forest matrix (Bobrowiec and Gribel, 2010).

6. Broader consequences of fragmentation

6.1. Ecological distortions are common

Many ecological interactions are altered in fragmented forests. Fragmented communities can pass through unstable transitional states that may not otherwise occur in nature (Terborgh et al. 2001). Moreover, species at higher trophic levels, such as predators and parasites, are often more vulnerable to fragmentation than are herbivores, thereby altering the structure and functioning of food webs (Didham et al. 1998b, Terborgh et al. 2001).

BDFFP findings suggest that even forest fragments that are unhunted, unlogged, and unburned have reduced densities of key mammalian seed dispersers. As a result, seed dispersal for the endemic, mammal-dispersed tree *Duckeodendron cestroides* was far lower in fragments, with just ~5% of the number of seeds being dispersed >10 m away from parent trees than in intact forest (Cramer et al. 2007a). Leaf herbivory appears reduced in fragments, possibly because of lower immigration of insect herbivores (Fáveri et al. 2008). Dung beetles exhibit changes in biomass and guild structure in fragments (Radtke et al. 2008) that could alter rates of forest nutrient cycling and secondary seed dispersal (Klein 1989, Andresen 2003). Exotic Africanized honeybees, a generalist pollinator, are abundant in matrix and edge

habitats and can alter pollination distances and gene flow for some tree species (Dick 2001, Dick et al. 2003). A bewildering variety of ecological distortions can pervade fragmented habitats, and a challenge for conservation biologists is to identify those of greatest importance and generality.

6.2. Fragmentation affects much more than biodiversity

Habitat fragmentation affects far more than biodiversity and interactions among species; many ecosystem functions, including hydrology (see above) and biochemical cycling, are also being altered. Among the most important of these are fundamental changes in forest biomass and carbon storage.

Carbon storage in fragmented forests is affected by a suite of interrelated changes. Many trees die near forest edges (Laurance et al. 1997, 1998a), including an alarmingly high proportion of large (≥ 60 cm dbh) canopy and emergent trees that store much forest carbon (Laurance et al. 2000). Fast-growing pioneer trees and lianas that proliferate in fragments are smaller and have lower wood density (Fig. 5), and thereby sequester much less carbon, than do the mature-phase trees they replace (Laurance et al. 2001b, 2006a). Based on current rates of forest fragmentation, the edge-related loss of forest carbon storage might produce up to 150 million tons of atmospheric carbon emissions annually, above and beyond that from tropical deforestation per se (Laurance et al. 1998c). Such discharge would exceed the yearly carbon emissions of the entire United Kingdom. Note, however, that most of this emission is already counted in the existing estimates of the impact of Amazonian land-use change because the deforestation emission estimates use forest biomass values for undegraded forest (Fearnside 2000). Because most deforestation occurs by expansion of already-existing clearings, forest edges (with reduced biomass) are the first areas to be cleared. Only the annual increase in the total length of forest edges represents an addition. Improved emissions estimates, accounting for degradation by logging, fire and fragmentation, are a high priority.

In addition, biomass is being redistributed in fragmented forests. Less biomass is stored in large, densely wooded old-growth trees and more in fast-growing pioneer trees, disturbance-loving lianas, woody debris, and leaf litter (Sizer et al. 2000, Nascimento and Laurance 2004, Vasconcelos and Luizão 2004). Finally, carbon cycling accelerates. The large, mature-phase trees that predominate in intact forests can live for many centuries or even millennia (Chambers et al. 1998, Laurance et al. 2004), sequestering carbon for long periods of time. However, the residence time of carbon in early successional trees, vines, and necromass (wood debris, litter), which proliferate in fragments, is far shorter (Nascimento and Laurance 2004). Other biochemical cycles, such as those affecting key nutrients like phosphorus (Sizer et al. 2000) and calcium (Vasconcelos and Luizão 2004), may also be altered in fragmented forests, given the striking changes in biomass dynamics, hydrology, and thermal regimes they experience.

7. Predicting species responses to fragmentation

7.1. Species losses are highly nonrandom

Local extinctions of species in the BDFFP fragments have occurred in a largely predictable sequence, with certain species being consistently more vulnerable than others. Among birds, a number of understory insectivores, including army ant-followers, solitary species, terrestrial foragers, and obligate mixed-flock members, are most susceptible to fragmentation. Others, including edge/ gap species, insectivores that use mixed flocks facultatively, hummingbirds, and many frugivores, are far less vulnerable (Antongiovanni and Metzger 2005, Stouffer et al. 2006, 2008). In a similar vein, among bats, gleaning predators are consistently the most vulnerable species whereas many frugivores respond positively to fragmentation and disturbance (Sampaio 2000, Bobrowiec and Gribel 2010, Rocha et al. 2013). Primates exhibit

similarly predictable patterns of species loss, with wide-ranging frugivores, especially the black spider-monkey, being most vulnerable (Boyle and Smith 2010a). Local extinctions in fragments follow a foreseeable pattern, with species assemblages in smaller fragments rapidly forming a nested subset of those in larger fragments (Stouffer et al. 2008). Random demographic and genetic processes may help to drive tiny populations into oblivion, but the species that reach this precarious threshold are far from random.

7.2. Fragmented communities are not neutral

An important corollary of nonrandom species loss is that fragmented forests are not neutral. Neutral theory (Hubbell 2001) assumes that species in diverse, space-limited communities, such as tropical trees, are competitively equivalent in order to make predictions about phenomena such as species-area curves, the relative abundances of species in communities, and the rate of species turnover in space. Hubbell (2001) emphasizes the potential relevance of neutral theory for predicting community responses to habitat fragmentation: for isolated communities, locally abundant species should be least extinction prone, with rare species being lost more frequently from random demographic processes. Over time, fragments should become dominated by initially abundant species, with rare species gradually vanishing; other ecological traits of species are considered unimportant.

Gilbert et al. (2006) tested the efficacy of neutral theory for predicting changes in tree communities at the BDFFP. Neutral theory effectively predicted the rate of local extinctions of species from plots in fragmented and intact forest as a function of the local diversity and the mortality rate of trees. However, in most fragments, the observed rate of change in species composition was 2–6 times faster than predicted by the theory. Moreover, the theory was wildly erroneous in predicting which species are most prone to local extinction. Rather than becoming increasingly dominated by initially common species, fragments in the BDFFP landscape have experienced striking increases in disturbance-loving pioneer species (Fig. 6) (Laurance et al. 2006a), which were initially rare when the fragments were created. As a model for predicting community responses to habitat fragmentation, neutral theory clearly failed, demonstrating that ecological differences among species strongly influence their responses to fragmentation.

7.3. Matrix use and area needs determine animal vulnerability

The responses of animal species to fragmentation appear largely governed by two key sets of traits. The first is their spatial requirements for forest habitat. In birds (Van Houtan et al. 2007) and mammals (Timo 2003), wide-ranging forest species are more vulnerable than are those with localized ranges and movements. Species with limited spatial needs, such as many small mammals (Malcolm 1997), hummingbirds (Stouffer et al. 2008), frogs (Tocher et al. 1997), and ants (Carvalho and Vasconcelos 1999), are generally less susceptible to fragmentation.

The second key trait for fauna is their tolerance of matrix habitats (Gascon et al. 1999), which comprises cattle pastures and regrowth forest in the BDFFP landscape. Populations of species that avoid the matrix will be entirely isolated in fragments, and therefore vulnerable to local extinction, whereas those that tolerate or exploit the matrix often persist (Laurance 1991, Malcolm 1997, Antongiovanni and Metzger 2005, Ferraz et al. 2007, Bobrowiec and Gribel 2010). At least among terrestrial vertebrates, matrix use is positively associated with tolerance of edge habitats (Laurance 2004b, Farneda 2013), an ability to traverse small clearings (Laurance et al. 2004, Laurance and Gomez 2005), and behavioral flexibility (Neckel-Oliveira and Gascon 2006, Stouffer et al. 2006, Van Houtan et al. 2006, Boyle and Smith 2010b). Within particular animal groups, such as beetles or small mammals,

traits such as body size and natural abundance are poor or inconsistent predictors of vulnerability (Laurance 1991, Didham et al. 1998a, Jorge 2008, Boyle and Smith 2010a). Natural abundance, however, is an important predictor of sensitivity to fragmentation for bats at the BDFFP (Farneda, 2013).

7.4. Disturbance tolerance and mutualisms affect plant vulnerability

Among plants, a different suite of factors is associated with vulnerability to fragmentation. Because fragments suffer chronically elevated tree mortality, faster-growing pioneer trees and lianas that favor treefall gaps are favored at the expense of slower-growing mature-phase trees (Laurance et al. 2006a, b). Pioneer species often flourish in the matrix and produce abundant small fruits that are carried into fragments by frugivorous birds and bats that move between the matrix and nearby fragments (Sampaio 2000, Nascimento et al. 2006). Especially vulnerable in fragments are the diverse assemblages of smaller subcanopy trees that are physiologically specialized for growth and reproduction in dark, humid, forest-interior conditions (Laurance et al. 2006b). Tree species that have obligate outbreeding systems, rely on animal seed dispersers, or have relatively large, mammal-dispersed seeds also appear vulnerable (Laurance et al. 2006b, Cramer et al. 2007b).

These combinations of traits suggest that plant communities in fragmented forests are structured primarily by chronic disturbances and microclimatic stresses and possibly also by alterations in animal pollinator and seed-disperser communities. For long-lived plants such as *Heliconia* species and many mature-phase trees, demographic models suggest that factors that reduce adult survival and growth—such as recurring wind disturbance and edge-related microclimatic stresses—exert a strong influence on population growth (Bruna 2003, Bruna and Oli 2005).

Differential tolerance to drought also seems to play a role on secondary forests. We find higher and significant mortality and lower biomass accumulation rates in *Cecropia*-dominated secondary forests, associated with drier years, while *Vismia*-dominated regrowth showed a non-significant, but similar trend. It is likely that different species assemblages account for the differential ability of these successional pathways to tolerate extreme climate events (Mesquita et al. 2012).

8. Broad perspectives

8.1. Long-term research is crucial

Many insights from the BDFFP would have been impossible in a shorter-term study. The exceptional vulnerability of large trees to fragmentation (Laurance et al. 2000) only became apparent after two decades of fragment isolation. Likewise, the importance of ephemeral events such as El Niño droughts (Williamson et al., 2000, Laurance et al. 2001c) and major windstorms (Laurance et al. 2007) would not have been captured in a less-enduring project. Many other key phenomena, such as the kinetics of species loss in fragments (Ferraz et al. 2003), the strong effects of matrix dynamics on fragmented bird and bat assemblages (Antongiovanni and Metzger 2005, Stouffer et al. 2006, Meyer et al. 2013), the divergence of fragments in different landscapes (Laurance et al. 2007), and the effects of fragmentation on rare or long-lived species (Benítez-Malvido and Martinez-Ramos 2003b, Ferraz et al., 2007), are only becoming understood after decades of effort.

Far more remains to be learned. For example, forest-simulation models parameterized with BDFFP data suggest that even small (<10 ha) fragments will require a century or more to stabilize in floristic composition and carbon storage (Groeneveld et al. 2009), given the long-lived nature of many tropical trees. Eventually, these fragments might experience a fundamental reorganization of their plant communities, given striking shifts in the composition of their tree, palm, liana, and herb seedlings (Scariot 2001; Benítez-Malvido and

Martinez-Ramos 2003a, Brum et al. 2008). If these newly recruited plants represent the future of the forest, then the BDFFP fragments will eventually experience dramatic changes in floristic composition—comparable to those observed in some other long-fragmented ecosystems (e.g. da Silva and Tabarelli 2000, Girão et al. 2007, Santos et al. 2010).

8.2. The BDFFP is a best-case scenario

Although forest fragments in the BDFFP are experiencing a wide array of ecological changes, it is important to emphasize that it is a controlled experiment. The fragments are square, not irregular, in shape. They are isolated by distances of just 80–650 m from large tracts of surrounding mature forest. They are embedded in a relatively benign matrix increasingly dominated by regrowth forest. And they lack many of the ancillary threats, such as selective logging, wildfires, and overhunting, that plague many fragmented landscapes and wildlife elsewhere in the tropics (e.g. Moura et al. 2014). Such threats can interact additively or synergistically with fragmentation, creating even greater perils for the rainforest biota (Laurance and Cochrane 2001, Michalski and Peres 2005, Brook et al. 2008). For these reasons, results from the BDFFP are clearly optimistic relative to many human-dominated landscapes elsewhere in the tropics.

9. Conservation lessons from the BDFFP

9.1. Amazonian reserves should be large and numerous

A key conclusion from BDFFP research is that nature reserves in Amazonia should ideally be very large—on the order of thousands to tens of thousands of square kilometers (Laurance 2005, Peres 2005). Only at this size will they be likely to maintain natural ecological processes and sustain viable populations of the many rare and patchily distributed species in the region (Ferraz et al. 2007, Radtke et al. 2008); provide resilience from rare calamities such as droughts and intense storms (Laurance et al. 2007); facilitate persistence of terrestrial and aquatic animals that migrate seasonally (Bührnheim and Fernandes 2003); buffer the reserve from large-scale edge effects including fires, forest desiccation, and human encroachment (Cochrane and Laurance 2002, Briant et al. 2010); maximize forest carbon storage (Laurance et al. 1997, 1998c); and provide resilience from future climatic and atmospheric changes—the effects of which are difficult to predict for Amazonia (Laurance and Useche 2009). Further, on the ancient soils of Central and Eastern Amazonia, low plant productivity translates into low population densities of many animals up the food chain, so reserves must be proportionately larger to harbor viable populations (Radtke et al. 2008, Deichmann et al. 2011, 2012).

Nature reserves in Amazonia should also be numerous and stratified across major river basins and climatic and edaphic gradients, in order to preserve locally endemic species (Bierregaard et al. 2001, Laurance, 2007). Further, the core areas of reserves should ideally be free of roads, which can promote human encroachment and hunting, internally fragment wildlife populations, and facilitate invasions of exotic species and fire (Laurance et al. 2009b).

9.2. Protect and reconnect fragments

Few landscapes are as intact as those in the Amazon. Around the world, biodiversity hotspots, which sustain the majority of species at risk of extinction, have, by definition, lost over 80% of their natural vegetation and what remains is typically in small fragments (Myers et al. 2000). The BDFFP makes recommendations here, too. Reconnecting isolated fragments by forest restoration will be an effective way of creating areas large enough to slow the rate of local species extinctions (Lima and Gascon 1999, Pimm and Jenkins 2005).

In such heavily fragmented landscapes, protecting remaining forest remnants is highly

desirable, as they are likely to be key sources of plant propagules and animal seed dispersers and pollinators (Mesquita et al. 2001, Chazdon et al. 2008). They may also act as stepping stones for animal movements (Laurance and Bierregaard 1997, Dick et al. 2003). In regions where forest loss is severe, forest fragments could also sustain the last surviving populations of locally endemic species, thereby underscoring their potential value for nature conservation (Arroyo-Rodríguez et al. 2009).

9.3. Fragmented landscapes can recover

A further lesson is that fragmented landscapes, if protected from fires and other major disturbances, can begin to recover in just a decade or two. Forest edges tend to ‘seal’ themselves, reducing the intensity of deleterious edge effects (Camargo and Kapos 1995, Didham and Lawton 1999, Mesquita et al. 1999). Secondary forests can develop quickly in the surrounding matrix (Mesquita et al. 2001), especially if soils and seedbanks are not depleted by overgrazing or repeated burning (Ribeiro et al. 2009, Norden et al. 2011). Secondary forests facilitate movements of many animal species (Gascon et al. 1999), allowing them to recolonize fragments from which they had formerly disappeared (Becker et al. 1991, Quintero and Roslin 2005, Stouffer et al. 2008, Bobrowiec and Gribel 2010, Boyle and Smith 2010a, Meyer et al. 2013). Species clinging to survival in fragments can also be rescued from local extinction via the genetic and demographic contributions of immigrants (Zartman and Nascimento 2006, Stouffer et al. 2008).

10. The future of the BDFFP

The BDFFP is one of the most enduring and influential ecological research projects in existence today (Gardner et al. 2009, Peres et al. 2010). From the prism of understanding habitat fragmentation, there are vital justifications for continuing it. The project, moreover, is engaged in far more than fragmentation research: it plays a leading role in training Amazonian scientists and decision-makers, and sustains long-term research on global-change phenomena, forest regeneration, and basic ecological studies.

In its 36-year history, the BDFFP has faced myriad challenges. These include, among others, the continuing fluctuations in currencies, challenges in obtaining research visas for foreign students and scientists, inadequate core funding from its US and Brazilian sponsors, and the vagaries of finding soft money for long-term research and to sustain a minimal number of workers to support infrastructure and logistics. Yet today the BDFFP faces a far more direct threat: encroachment from colonists and hunters. Since the late 1990s, the paving of the 1100-km-long Manaus–Venezuela (BR-174) highway has greatly accelerated forest colonization and logging north of the city. SUFRAMA, a Brazilian federal agency that controls an expanse of land north of Manaus that includes the BDFFP, has begun settling families in farming plots around the immediate periphery of the study area. At least six colonization projects involving 180 families are planned for the area (Laurance and Luizão 2007). These settlements could be the beginning of a dramatic influx into the area, especially if the proposed BR-319 highway between Manaus and Rondônia, a major deforestation hotspot in southern Amazonia, is completed as planned (Fearnside and Graça 2006).

To date, BDFFP staff and supporters have managed to stave off most of the colonization projects—which also threaten to bisect the Central Amazonian Conservation Corridor, a budding network of protected and indigenous lands that is one of the most important conservation areas in the entire Amazon basin (Laurance and Luizão 2007). Yet it is an uphill battle against a government bureaucracy that appears myopically determined to push ahead with colonization at any cost—despite the fact that colonists can barely eke out a living on the region’s infamously poor soils (Fearnside and Leal Filho, 2001). That such a globally important research project and conservation area could be lost seems unthinkable.

That it could be lost for such a limited gain seems tragic.

Amazon forest is under stress from a variety of global changes that are expected to increase in the coming decades. Beyond the considerable contributions of the BDFFP to date in providing information relevant to understanding these changes, the project is uniquely well placed to track the impacts of these changes as they occur. The BDFFP must continue its role in contributing to the scientific basis for more serious global efforts to contain the current human destruction of the environment at both the global and regional level.

Acknowledgements

We thank Laszlo Nagy and an anonymous referee for helpful comments on the manuscript. The National Institute for Amazonian Research (INPA), Smithsonian Institution, US National Science Foundation, Brazilian Science Foundation (CNPq), Amazonian State Science Foundation (FAPEAM), NASA-LBA program, USAID, Mellon Foundation, Blue Moon Fund, Marisla Foundation, and other organizations generously supported the BDFFP. Substantial parts of this text are updated from Laurance et al. (2011). This is publication number 640 in the BDFFP technical series.

Box 1. The Biological Dynamics of Forest Fragments Project

Since its inception in 1979, the Biological Dynamics of Forest Fragments Project (BDFFP) has been assessing the impacts of fragmentation on the Amazon rainforest and biota (Lovejoy et al. 1986, Bierregaard et al. 1992, Pimm 1998, Laurance et al. 2002, 2011). Today, it is the world's largest and longest-running experimental study of habitat fragmentation, as well as one of the most highly cited ecological investigations ever conducted (Gardner et al. 2009, Peres et al. 2010, Pitman et al. 2011). The BDFFP has also been a global leader in research, training, and capacity development, with over 640 publications (<http://pdbff.inpa.gov.br>), more than 180 student theses, over 700 graduate students and conservation professionals participating in sponsored courses, and over 1000 student interns to date.

The BDFFP is located 80 km north of Manaus, Brazil and spans ~1000 km². The topography is relatively flat (80–160 m elevation) but dissected by numerous stream gullies. The heavily weathered, nutrient-poor soils of the study area are typical of large expanses of the Amazon Basin. Rainfall ranges from 1900 to 3500 mm annually with a moderately strong dry season from June to October. The forest canopy is 30–37 m tall, with emergent trees to 55 m. Species richness of trees (≥ 10 cm DBH) often exceeds 280 species ha⁻¹ (Oliveira and Mori 1999, Laurance et al. 2010b) with a comparably high level of diversity also evident in many other plant and animal taxa.

The study area includes three large cattle ranges (~5000 ha each) containing 11 forest fragments (five of 1 ha, four of 10 ha, and two of 100 ha), and expanses of nearby continuous forest that serve as experimental controls. In the early 1980s, the fragments were isolated from nearby intact forest by distances of 80–650 m by clearing and burning the surrounding forest. A key feature was that pre-fragmentation censuses were conducted for many animal and plant groups (e.g. trees, understory birds, small mammals, primates, frogs, many invertebrate taxa), thereby allowing long-term changes in these groups to be assessed far more confidently than in most other fragmentation studies.

Because of poor soils and low productivity, the ranches surrounding the BDFFP fragments were largely abandoned after government fiscal incentives dried up from 1984 onwards. Secondary forests (initially dominated by *Vismia* spp in areas that were cleared and burned, or by *Cecropia* spp in areas that were cleared without fire) proliferated in many formerly forested areas (Mesquita et al. 2001). Some of the regenerating areas initially dominated by *Cecropia* spp later developed into quite mature (>20 m tall), species-rich

secondary forests. *Vismia*-dominated regrowth, which is relatively species poor, is changing far more slowly (Norden et al. 2011, Williamson et al. 2014). To help maintain isolation of the experimental fragments, 100 m-wide strips of regrowth were cleared and burned around each fragment on 4–5 occasions, most recently in 2013–2014. Additional human disturbances that harm many fragmented landscapes in the Amazon, such as major fires and logging, are largely prevented at the BDFFP. Hunting pressure has been very limited until recently, following a government decision to increase colonization in the general area (Laurance and Luizão 2007). Laurance and Bierregaard (1997) and Bierregaard et al. (2001) provide detailed descriptions of the study area and design.

References

- Andresen E (2003) Effect of forest fragmentation on dung beetle communities and functional consequences for plant regeneration. *Ecography* 26:87–97.
- Antongiovanni M, Metzger JP (2005) Influence of matrix habitats on the occurrence of insectivorous bird species in Amazonian forest fragments. *Biol Conserv* 122:441–451.
- Arroyo-Rodríguez V, Pineda E, Escobar F, Benítez-Malvido J (2009) Conservation value of small patches to plant species diversity in highly fragmented landscapes. *Conserv Biol* 23:729–739.
- Asner GP, Knapp D, Broadbent E, Oliveira P, Keller M, Silva J (2005) Selective logging in the Brazilian Amazon. *Science* 310:480–482.
- Avissar R, Liu Y (1996) A three-dimensional numerical study of shallow convective clouds and precipitation induced by land-surface forcing. *J Geophys Res* 101:7499–7518.
- Avissar R, Schmidt T (1998) An evaluation of the scale at which ground-surface heat flux patchiness affects the convective boundary layer using a large-eddy simulation model. *J Atmos Sci* 55:2666–2689.
- Baker TR, Phillips OL, Malhi Y, Almeida S, Arroyo L, Di Fiore A, et al. (2004) Increasing biomass in Amazonian forest plots. *Phil Trans Roy Soc B* 359:353–365.
- Barlow J, Peres CA, Henriques L, Stouffer PC, Wunderle J (2006) The responses of understorey birds to forest fragmentation, logging and wildfires: an Amazonian synthesis. *Biol Conserv* 128:182–192.
- Becker P, Moure JB, Peralta F (1991) More about euglossine bees in Amazonian forest fragments. *Biotropica* 23:586–591.
- Benítez-Malvido J (1998) Impact of forest fragmentation on seedling abundance in a tropical rain forest. *Conserv Biol* 12:380–389.
- Benítez-Malvido J, Martinez-Ramos M (2003a) Influence of edge exposure on tree seedling species recruitment in tropical rain forest fragments. *Biotropica* 35:530–541.
- Benítez-Malvido J, Martinez-Ramos M (2003b) Impact of forest fragmentation on understory plant species richness in Amazonia. *Conserv Biol* 17:389–400.
- Bierregaard RO, Lovejoy TE, Kapos V, dos Santos AA, Hutchings RW (1992) The biological dynamics of tropical rainforest fragments. *Bioscience* 42:859–866.
- Bierregaard R.O., Gascon C., Lovejoy T.E., Mesquita R. (Eds), 2001. *Lessons from Amazonia: Ecology and Conservation of a Fragmented Forest*. Yale University Press, New Haven, Connecticut.
- Bobrowiec PED, Gribel R (2010) Effects of different secondary vegetation types on bat community composition in Central Amazonia, Brazil. *Anim Conserv* 13:204–216.
- Bohlman S, Laurance WF, Laurance SG, Nascimento H, Fearnside PM, Andrade A (2008) Effects of soils, topography, and geographic distance in structuring central Amazonian tree communities. *J Veg Sci* 19:863–874.
- Boyle SA, Smith AT (2010a) Can landscape and species characteristics predict primate presence in forest fragments in the Brazilian Amazon? *Biol Conserv* 143:1134–1143.

- Boyle SA, Smith AT (2010b) Behavioral modifications in northern bearded saki monkeys (*Chiropotes satanas chiropotes*) in forest fragments of central Amazonia. *Primates* 51:43–51.
- Brazil, RADAMBRASIL (1973-1983) Levantamento de Recursos Naturais. Ministério das Minas e Energia, Departamento Nacional de Produção Mineral, Rio de Janeiro, Brazil.
- Briant G, Gond V, Laurance SGW (2010) Habitat fragmentation and the desiccation of forest canopies: a case study from eastern Amazonia. *Biol Conserv* 143:2763–2769.
- Broadbent E, Asner GP, Keller M, Knapp D, Oliveira P, Silva J (2008) Forest fragmentation and edge effects from deforestation and selective logging in the Brazilian Amazon. *Biol Conserv* 140:142–155.
- Brook BW, Sodhi NS, Bradshaw CJA (2008) Synergisms among extinction drivers under global change. *Trends Ecol Evol* 23:453–460.
- Brown KS, Hutchings RW (1997) Disturbance, fragmentation, and the dynamics of diversity in Amazonian forest butterflies. In: Laurance WF, Bierregaard RO (Eds), *Tropical forest remnants: Ecology, management, and conservation of fragmented communities*. University of Chicago Press, Chicago, pp 91–110.
- Brum HD, Nascimento H, Laurance WF, Andrade A, Laurance SGW, Luizão R (2008) Rainforest fragmentation and the demography of the economically important palm *Oenocarpus bacaba* in central Amazonia. *Plant Ecol* 199:209–215.
- Bruna EM (1999) Seed germination in rainforest fragments. *Nature* 402:139.
- Bruna EM (2003) Are plant populations in fragmented habitats recruitment limited? Tests with an Amazonian herb. *Ecology* 84:932–947.
- Bruna E, Oli M (2005) Demographic consequences of habitat fragmentation for an Amazonian understory plant: analysis of life-table response experiments. *Ecology* 86:1816–1824.
- Bruna EM, Vasconcelos HL, Heredia S (2005) The effect of habitat fragmentation on communities of mutualists: a test with Amazonian ants and their host plants. *Biol Conserv* 124:209–216.
- Bührnheim CM, Fernandes CC (2003) Structure of fish assemblages in Amazonian rainforest streams: effects of habitats and locality. *Copeia* 2003:255–262.
- Butler RA, Laurance WF (2009) Is oil palm the next emerging threat to the Amazon? *Trop Cons Sci* 2:1-10.
- Camargo JLC, Kapos V (1995) Complex edge effects on soil moisture and microclimate in central Amazonian forests. *J Trop Ecol* 11:205–211.
- Carvalho KS, Vasconcelos HL (1999) Forest fragmentation in central Amazonia and its effects on litter-dwelling ants. *Biol Conserv* 91:151–158.
- Chambers JQ, Higuchi N, Schimel JP (1998) Ancient trees in Amazonia. *Nature* 391:135–136.
- Chave J, Muller-Landau, HC, Baker TR, Easdale TA, ter Steege H, Webb CO (2006) Regional and phylogenetic variation of wood density across 2,456 neotropical tree species. *Ecol Applic* 16:2356-2367.
- Chazdon RL, Harvey CA, Komar O, Griffith DM, Ferguson BG, Martinez- Ramos M, Morales H, Nigh R, Soto-Pinto L, van Breugel M, Philpott SM (2008) Beyond reserves: a research agenda for conserving biodiversity in human-modified tropical landscapes. *Biotropica* 41:142–153.
- Cochrane MA, Laurance WF (2002) Fire as a large-scale edge effect in Amazonian forests. *J Trop Ecol* 18:311–325.
- Cochrane MA, Laurance WF (2008) Synergisms among fire, land use, and climate change in the Amazon. *Ambio* 37:522–527.

- 841 Condon MA, Sasek TW, Strain BR (1992) Allocation patterns in two tropical vines in
842 response to increased atmospheric CO₂. *Funct Ecol* 6:680-685.
- 843 Cox PM, Betts RA, Collins M, Harris PP, Huntingford C, Jones CD (2004) Amazonian forest
844 dieback under climate-carbon cycle projections for the 21st century. *Theoret Appl*
845 *Climatol* 78:137-156.
- 846 Cox PM, Betts RA, Jones CD, Spall SA, Totterdell IJ (2000) Acceleration of global warming
847 due to carbon cycle feedbacks in a coupled climate model. *Nature* 408:184-187
- 848 Cox PM, Pearson D, Booth BB, Friedlingstein P, Huntingford C, Jones CD, Luke CMD
849 (2013) Sensitivity of tropical carbon to climate change constrained by carbon dioxide
850 variability. *Nature* 494:341-344.
- 851 Cramer JM, Mesquita R, Bentos T, Moser B, Williamson GB (2007a) Forest fragmentation
852 reduces seed dispersal of *Duckeodendron cestroides*, a Central Amazon endemic.
853 *Biotropica* 39:709-718.
- 854 Cramer JM, Mesquita R, Williamson GB (2007b) Forest fragmentation differentially affects
855 seed dispersal of large and small-seeded tropical trees. *Biol Conserv* 137:415-423.
- 856 Dai A (2012) Increasing drought under global warming in observations and models. *Nature*
857 *Climate Change* 3:52-58.
- 858 D'Angelo S, Andrade A, Laurance SG, Laurance WF, Mesquita R (2004) Inferred causes of
859 tree mortality in fragmented and intact Amazonian forests. *J Trop Ecol* 20:243-246.
- 860 da Silva JMC, Tabarelli M (2000) Tree species impoverishment and the future flora of the
861 Atlantic forest of northeast Brazil. *Nature* 404:72-74.
- 862 Develey P, Stouffer PC (2001) Roads affect movements by understory mixed-species flocks
863 in central Amazonian Brazil. *Conserv Biol* 15:1416-1422.
- 864 Deichmann JL, Lima A, Williamson GB (2011) Effects of geomorphology and primary
865 productivity on Amazonian leaf litter herpetofauna. *Biotropica* 43:149-156.
- 866 Deichmann JL, Toft CA, Deichmann PM, Lima AP, Williamson GB (2013) Neotropical
867 primary productivity affects biomass of leaf-litter herpetofaunal assemblage. *J Trop*
868 *Ecol* 28:427-435.
- 869 Diamond JM, Bishop KD, Balen SV (1987) Bird survival in an isolated Javan woodland:
870 island or mirror? *Conserv Biol* 1:132-142.
- 871 Dick CW (2001) Genetic rescue of remnant tropical trees by an alien pollinator. *Proc Roy Soc*
872 *B* 268:2391-2396.
- 873 Dick CW, Etchelecu G, Austerlitz F (2003) Pollen dispersal of tropical trees (*Dinizia excelsa*:
874 *Fabaceae*) by native insects and African honeybees in pristine and fragmented
875 Amazonian rainforest. *Mol Ecol* 12:753-764.
- 876 Didham RK, Hammond PM, Lawton JH, Eggleton P, Stork NE (1998a) Beetle species
877 responses to tropical forest fragmentation. *Ecol Monogr* 68:295-303.
- 878 Didham RK, Lawton JH (1999) Edge structure determines the magnitude of changes in
879 microclimate and vegetation structure in tropical forest fragments. *Biotropica* 31:17-
880 30.
- 881 Didham RK, Lawton JH, Hammond PM, Eggleton P (1998b) Trophic structure stability and
882 extinction dynamics of beetles (Coleoptera) in tropical forest fragments. *Proc Roy Soc*
883 *B* 353:437-451.
- 884 Farneda FZ (2013) Bat species vulnerability to forest fragmentation in the Central Amazon.
885 M.Sc. Thesis. University of Lisbon, Portugal.
- 886 Fáveri SB, Vasconcelos HL, Dirzo R (2008) Effects of Amazonian forest fragmentation on
887 the interaction between plants, insect herbivores, and their natural enemies. *J Trop*
888 *Ecol* 24:57-64.
- 889 Fearnside PM (1997) Wood density for estimating forest biomass in Brazilian Amazonia. *For*
890 *Ecol Manage* 90:59-89. |

- Fearnside PM (2000) Global warming and tropical land-use change: Greenhouse gas emissions from biomass burning, decomposition and soils in forest conversion, shifting cultivation and secondary vegetation. *Climatic Change* 46:115-158.
- Fearnside PM (2001a) Soybean cultivation as a threat to the environment in Brazil. *Environ Conserv* 28:23–38.
- Fearnside PM (2001b) Saving tropical forests as a global warming countermeasure: An issue that divides the environmental movement. *Ecol Econ* 39:167-184.
- Fearnside PM (2002) *Avança Brasil: Environmental and social consequences of Brazil's planned infrastructure in Amazonia*. *Environ Manage* 30:748-763.
- Fearnside PM (2007) Brazil's Cuiabá-Santarém (BR-163) Highway: The environmental cost of paving a soybean corridor through the Amazon. *Environ Manage* 39:601-614.
- Fearnside PM (2012) Brazil's Amazon forest in mitigating global warming: Unresolved controversies. *Climate Policy* 12:70-81.
- Fearnside PM (2013) Vines, CO₂ and Amazon forest dieback. *Nature Online Comment* (<http://www.nature.com/nature/journal/vaop/ncurrent/full/nature11882.html>).
- Fearnside, PM (2014) The impact of land use on carbon stocks and fluxes. (this volume).
- Fearnside PM, Graça PMLA (2006) BR-319: Brazil's Manaus–Porto Velho highway and the potential impact of linking the arc of deforestation to central Amazonia. *Environ Manage* 38:705–716.
- Fearnside PM, Leal Filho N (2001) Soil and development in Amazonia: Lessons from the Biological Dynamics of Forest Fragments Project. In: Bierregaard RO, Gascon C, Lovejoy TE, Mesquita R (Eds) *Lessons from Amazonia: The ecology and conservation of a fragmented forest*. Yale University Press, New Haven, Connecticut, USA, pp 291-312.
- Ferraz G, Nichols JD, Hines J, Stouffer PC, Bierregaard RO, Lovejoy TE (2007) A large-scale deforestation experiment: effects of patch area and isolation on Amazon birds. *Science* 315:238–241.
- Ferraz G, Russell GJ, Stouffer PC, Bierregaard RO, Pimm SL, Lovejoy TE (2003) Rates of species loss from Amazonian forest fragments. *Proc Nat Acad Sci USA* 100:14069–14073.
- Finer M, Jenkins C, Pimm SL, Keane B, Ross C (2008) Oil and gas projects in the western Amazon: threats to wilderness, biodiversity, and indigenous peoples. *PLoS ONE* 3:e2932.
- Fowler HG, Silva CA, Ventincinque E (1993) Size, taxonomic and biomass distributions of flying insects in central Amazonia: forest edge vs. understory. *Rev Biol Trop* 41:755–760.
- Gardner TA, Barlow J, Chazdon R, Ewers R, Harvey C, Peres CA, Sodhi NS (2009) Prospects for tropical forest biodiversity in a human-modified world. *Ecol Lett* 12:561–582.
- Gascon C (1993) Breeding habitat use by Amazonian primary-forest frog species at the forest edge. *Biodiv. Conserv.* 2:438–444.
- Gascon C, Lovejoy TE, Bierregaard RO, Malcolm JR, Stouffer PC, Vasconcelos H, Laurance WF, Zimmerman B, Tocher M, Borges S (1999) Matrix habitat and species persistence in tropical forest remnants. *Biol Conserv* 91:223–229.
- Gibbs HK, Reusch AS, Achard F, Clayton MK, Holmgren P, Ramankutty N, Foley JA (2010) Tropical forests were the primary sources of new agricultural lands in the 1980s and 1990s. *Proc Nat Acad Sci USA* 107:16732–16737.
- Gibson L, Lee TM, Koh LP, Brook BW, Gardner, TA, Barlow J, Peres CA, Bradshaw CJA, Laurance WF, Lovejoy TE, Sodhi NS (2011) Primary forests are irreplaceable for sustaining tropical biodiversity. *Nature* 478:378-381.

- Gilbert B, Laurance WF, Leigh EG, Nascimento H (2006) Can neutral theory predict the responses of Amazonian tree communities to forest fragmentation? *Am Nat* 168:304–317.
- Gilbert KA, Setz EZF (2001) Primates in a fragmented landscape: six species in central Amazonia. In: Laurance WF, Bierregaard RO (Eds) *Tropical forest remnants: Ecology, management and conservation of fragmented communities*. University of Chicago Press, Chicago, pp 207–221.
- Girão LC, Lopes AV, Tabarelli M, Bruna EM (2007) Changes in tree reproductive traits reduce functional diversity in a fragmented Atlantic forest landscape. *PLoS ONE* 2:e908.
- Good P, Jones CD, Lowe JA, Betts RA, Gedney N (2013) Comparing tropical forest projections from two generations of Hadley Centre Earth System models, HadGEM2-ES and HadCM3LC. *J. Climate* 26:495–511.
- Groeneveld J, Alves L, Bernacci L, Catharino E, Knogge C, Metzger J, Pütz S, Huth A (2009) The impact of fragmentation and density regulation on forest succession in the Atlantic rain forest. *Ecol Model* 220:2450–2459.
- Harper LH (1989) The persistence of ant-following birds in small Amazonian forest fragments. *Acta Amazonica* 19:249–263.
- Hubbell SP (2001) *The Neutral Theory of Biodiversity and Biogeography*. Princeton University Press, Princeton, New Jersey.
- Huntingford C, Zelazowski P, Galbraith D, Mercado LM, Sitch S, Fisher R, Lomas M, Walker AP, Jones CD, Booth BB, Malhi Y, Hemming D, Kay G, Good P, Lewis SL, Phillips OL, Atkin OK, Lloyd J, Gloor E, Zaragoza-Castells J, Meir P, Betts R, Harris PP, Nobre C, Marengo J, Cox PM (2013) Simulated resilience of tropical rainforests to CO₂-induced climate change. *Nature Geosci.* 6:268–273.
- Ingwell LL, Wright SJ, Becklund KK, Hubbell SP, Schnitzer SA (2010) The impact of lianas on 10 years of tree growth and mortality on Barro Colorado Island, Panama. *J Ecol* 98:879–887.
- IPCC (2013) Summary for policymakers. In: Stocker T F et al. (Eds) *Climate change 2013: The physical science basis*. Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, UK.
- Janzen DH (1983) No park is an island: Increase in interference from outside as park size increases. *Oikos* 41:402–410.
- Jorge ML (2008) Effects of forest fragmentation on two sister genera of Amazonian rodents (*Myoprocta acouchy* and *Dasyprocta leporina*). *Biol Conserv* 141:617– 623.
- Kapos V (1989) Effects of isolation on the water status of forest patches in the Brazilian Amazon. *J Trop Ecol* 5:173–185.
- Kapos V, Ganade G, Matusi E, Victoria RL (1993) Delta 13C as an indicator of edge effects in tropical rainforest reserves. *J Ecol* 81:425–432.
- Killeen TJ (2007) *A Perfect Storm in the Amazon Wilderness: Development and Conservation in the Context of the Initiative for the Integration of the Regional Infrastructure of South America (IIRSA)*. Conservation International, Washington, DC.
- Klein BC (1989) Effects of forest fragmentation on dung and carrion beetle communities in central Amazonia. *Ecology* 70:1715–1725.
- Laurance SGW (2004b) Responses of understory rain forest birds to road edges in central Amazonia. *Ecol Applic* 14:1344–1357.
- Laurance SGW, Andrade A, Laurance WF (2010a) Unanticipated effects of stand dynamism on Amazonian tree diversity. *Biotropica* 42:429–434.
- Laurance SGW, Gomez MS (2005) Clearing width and movements of understory rainforest

- 991 birds. *Biotropica* 37:149–152.
- 992 Laurance SGW, Laurance WF, Andrade A, Fearnside PM, Harms K, Luizão R, (2010b)
- 993 Influence of soils and topography on Amazonian tree diversity: a landscape-scale
- 994 study. *J Veg Sci* 21:96–106.
- 995 Laurance SGW, Laurance WF, Nascimento H, Andrade A, Fearnside PM, Rebello E, Condit
- 996 R (2009a) Long-term variation in Amazon forest dynamics. *J Veg Sci* 20:323–333.
- 997 Laurance SGW, Stouffer PC, Laurance WF (2004) Effects of road clearings on movement
- 998 patterns of understory rainforest birds in central Amazonia. *Conserv Biol* 18:1099–
- 999 1109.
- 1000 Laurance WF (1991) Ecological correlates of extinction proneness in Australian tropical
- 1001 rainforest mammals. *Conserv Biol* 5:79–89.
- 1002 Laurance WF (2001) The hyper-diverse flora of the central Amazon: an overview. In:
- 1003 Bierregaard RO, Gascon C, Lovejoy TE, Mesquita R (Eds), *Lessons from Amazonia:*
- 1004 *Ecology and conservation of a fragmented forest.* Yale University Press, New Haven,
- 1005 Connecticut, pp 47–53.
- 1006 Laurance WF (2002) Hyperdynamism in fragmented habitats. *J Veg Sci* 13:595–602.
- 1007 Laurance WF (2004a) Forest–climate interactions in fragmented tropical landscapes. *Phil.*
- 1008 *Trans. Roy. Soc. B* 359:345–352.
- 1009 Laurance WF (2005) When bigger is better: the need for Amazonian megareserves. *Trends*
- 1010 *Ecol Evol* 20:645–648.
- 1011 Laurance WF (2007) Have we overstated the tropical biodiversity crisis? *Trends Ecol Evol*
- 1012 22:65–70.
- 1013 Laurance WF (2008) Theory meets reality: how habitat fragmentation research has
- 1014 transcended island biogeographic theory. *Biol Conserv* 141:1731–1744.
- 1015 Laurance WF, Andrade A, Magrach, A, Camargo JLC, Valsko J, Campbell M, Fearnside PM,
- 1016 Edwards W, Lovejoy TE, Laurance SGW (2014a) Long-term changes in liana
- 1017 abundance and forest dynamics in undisturbed Amazonian forests. *Ecology* 95:1604–
- 1018 1611.
- 1019 Laurance WF, Andrade A, Magrach A, Camargo JLC, Campbell M, Fearnside PM, Edwards
- 1020 W, Valsko J, Lovejoy TE, Laurance SGW (2014b) Apparent environmental synergism
- 1021 drives the dynamics of Amazonian forest fragments. *Ecology* 95: 3018–3026.
- 1022 Laurance WF, Bierregaard RO (Eds) (1997) *Tropical forest remnants: Ecology, management,*
- 1023 *and conservation of fragmented communities.* University of Chicago Press, Chicago.
- 1024 Laurance WF, Camargo J, Luizão RCC, Laurance SGW, Pimm SL, Bruna E, Stouffer PC,
- 1025 Williamson GB, Benitez-Malvido J, Vasconcelos H, Van Houtan K, Zartman CE,
- 1026 Boyle S, Didham RK, Andrade A, Lovejoy TE (2011) The fate of Amazonian forest
- 1027 fragments: A 32-year investigation. *Biol Conserv* 144:56–67.
- 1028 Laurance WF, Cochrane MA (2001) Synergistic effects in fragmented landscapes. *Conserv*
- 1029 *Biol* 15:1488–1489.
- 1030 Laurance WF, Cochrane M, Bergen S, Fearnside PM, Delamonica P, Barber C, D’Angelo S,
- 1031 Fernandes T (2001a) The future of the Brazilian Amazon. *Science* 291:438–439.
- 1032 Laurance WF, Delamonica P, Laurance SGW, Vasconcelos HL, Lovejoy TE (2000)
- 1033 Rainforest fragmentation kills big trees. *Nature* 404:836.
- 1034 Laurance WF, Fearnside PM, Laurance SGW, Delamonica P, Lovejoy TE, Rankin-de Merona
- 1035 JM, Chambers JQ, Gascon C (1999) Relationship between soils and Amazon forest
- 1036 biomass: a landscape-scale study. *For Ecol Manage* 118:127–138.
- 1037 Laurance WF, Ferreira LV, Rankin-de Merona JM, Laurance SG (1998a) Rain forest
- 1038 fragmentation and the dynamics of Amazonian tree communities. *Ecology* 79:2032–
- 1039 2040.
- 1040 Laurance WF, Ferreira LV, Rankin-de Merona JM, Laurance SGW, Hutchings R, Lovejoy

- TE (1998b) Effects of forest fragmentation on recruitment patterns in Amazonian tree communities. *Conserv Biol* 12:460–464.
- Laurance WF, Goosem M, Laurance SGW (2009b) Impacts of roads and linear clearings on tropical forests. *Trends Ecol Evol* 24:659–669.
- Laurance WF, Laurance SGW, Delamonica P (1998c) Tropical forest fragmentation and greenhouse gas emissions. *For Ecol Manage* 110:173–180.
- Laurance WF, Laurance SGW, Ferreira LV, Rankin-de Merona J, Gascon C, Lovejoy TE (1997) Biomass collapse in Amazonian forest fragments. *Science* 278:1117–1118.
- Laurance WF, Lovejoy TE, Vasconcelos H, Bruna E, Didham R, Stouffer P, Gascon C, Bierregaard R, Laurance SG, Sampaio E (2002) Ecosystem decay of Amazonian forest fragments: a 22-year investigation. *Conserv Biol* 16:605–618.
- Laurance WF, Luizão RCC (2007) Driving a wedge into the Amazon. *Nature* 448:409–410.
- Laurance WF, Nascimento H, Laurance SG, Andrade A, Ewers R, Harms K, Luizão R, Ribeiro J (2007) Habitat fragmentation, variable edge effects, and the landscape-divergence hypothesis. *PLoS ONE* 2:e1017.
- Laurance WF, Nascimento H, Laurance SG, Andrade A, Fearnside PM, Ribeiro J (2006a) Rain forest fragmentation and the proliferation of successional trees. *Ecology* 87:469–482.
- Laurance WF, Nascimento H, Laurance SG, Andrade A, Ribeiro J, Giraldo JP, Lovejoy TE, Condit R, Chave J, D'Angelo S (2006b) Rapid decay of tree community composition in Amazonian forest fragments. *Proc Nat Acad Sci USA* 103:19010–19014.
- Laurance WF, Nascimento H, Laurance SG, Condit R, D'Angelo S, Andrade A (2004) Inferred longevity of Amazonian rainforest trees based on a long-term demographic study. *For Ecol Manage* 190:131–143.
- Laurance WF, Peres CA (Eds) (2006) *Emerging Threats to Tropical Forests*. University of Chicago Press, Chicago, USA.
- Laurance WF, Perez-Salicrup D, Delamonica P, Fearnside PM, D'Angelo S, Jerozolinski A, Pohl L, Lovejoy TE (2001b) Rain forest fragmentation and the structure of Amazonian liana communities. *Ecology* 82:105–116.
- Laurance WF, Useche DC (2009) Environmental synergisms and extinctions of tropical species. *Conserv Biol* 23:1427–1437.
- Laurance WF, Williamson GB (2001) Positive feedbacks among forest fragmentation, drought, and climate change in the Amazon. *Conserv Biol* 15:1529–1535.
- Laurance WF, Williamson GB, Delamonica P, Olivera A, Gascon C, Lovejoy TE, Pohl L (2001c) Effects of a strong drought on Amazonian forest fragments and edges. *J Trop Ecol* 17:771–785.
- Leidner AK, Haddad NM, Lovejoy TE (2010) Does tropical forest fragmentation increase long-term variability of butterfly communities? *PLoS ONE* 5:e9534.
- Lima M, Gascon C (1999) The conservation value of linear forest remnants in central Amazonia. *Biol Conserv* 91:241–247.
- Lovejoy TE, Bierregaard RO, Rylands AB, Malcolm JR, Quintela C, Harper L, Brown K, Powell A, Powell G, Schubart H, Hays M (1986) Edge and other effects of isolation on Amazon forest fragments. In: Soulé ME (Ed), *Conservation biology: The science of scarcity and diversity*. Sinauer, Sunderland, Massachusetts, pp 257–285.
- Lovejoy TE, Oren DC (1981) Minimum critical size of ecosystems. In: Burgess RL, Sharp DM (Eds), *Forest-island dynamics in man-dominated landscapes*. Springer-Verlag, New York, pp 7–12.
- Lovejoy TE, Rankin JM, Bierregaard RO, Brown KS, Emmons LH, Van der Voort ME (1984) Ecosystem decay of Amazon forest fragments. In: Nitecki MH (Ed), *Extinctions*. University of Chicago Press, Chicago, pp 295–325.

- 1091 Malcolm JR (1994) Edge effects in central Amazonian forest fragments. *Ecology* 75:2438–
1092 2445.
- 1093 Malcolm JR (1997) Biomass and diversity of small mammals in Amazonian forest fragments.
1094 In: Laurance WF, Bierregaard RO (Eds) *Tropical forest remnants: Ecology,*
1095 *management, and conservation of fragmented communities.* University of Chicago
1096 Press, Chicago, pp 207–221.
- 1097 Malcolm JR (1998) A model of conductive heat flow in forest edges and fragmented
1098 landscapes. *Clim Change* 39:487–502.
- 1099 Massoca PES, Jakovac ACC, Vizcarra TB, Williamson GB, Mesquita RCG (2012)
1100 Dinâmica e trajetórias da sucessão secundária na Amazônia central. *Bol Mus Para*
1101 *Emílio Goeldi Cienc Nat Belém* 7:235-250.
- 1102 Mesquita R, Delamônica P, Laurance WF (1999) Effects of surrounding vegetation on edge-
1103 related tree mortality in Amazonian forest fragments. *Biol Conserv* 91:129–134.
- 1104 Mesquita RCC, Bentos TV, Jakovac ACC, Massoca P, Williamson GB (2012) Amazonian
1105 secondary forests respond to climatic variation: biomass dynamics. 49th Annual
1106 Meeting of the Association for Tropical Biology and Conservation, Bonito, MS,
1107 Brazil.
- 1108 Mesquita R, Ickes K, Ganade G, Williamson GB (2001) Alternative successional pathways in
1109 the Amazon basin. *J Ecol* 89:528–537.
- 1110 Mestre LAM, Gasnier TR (2008) Populações de aranhas errantes do gênero *Ctenus* em
1111 fragmentos florestais na Amazônia Central. *Acta Amazonica* 38:159–164.
- 1112 Michalski F, Peres CA (2005) Anthropogenic determinants of primate and carnivore local
1113 extinctions in a fragmented forest landscape of Southern Amazonia. *Biol Conserv*
1114 124:383–396.
- 1115 Moura NG, Lees AC, Andretti CB, Davis BJW, Solar RRC, Aleixo A, Barlow J, Ferreira J,
1116 Gardner TA (2014) Avian biodiversity in multiple-use landscapes of the Brazilian
1117 Amazon. *Biol Conserv* 167:339-348.
- 1118 Myers N, Mittermeier RA, Mittermeier CG, Fonseca GAB, Kent J (2000) Biodiversity
1119 hotspots for conservation priorities. *Nature* 403:853–858.
- 1120 Nascimento H, Andrade A, Camargo J, Laurance WF, Laurance SG, Ribeiro J (2006) Effects
1121 of the surrounding matrix on tree recruitment in Amazonian forest fragments. *Conserv*
1122 *Biol* 20:853–860.
- 1123 Nascimento H, Laurance WF (2004) Biomass dynamics in Amazonian forest fragments. *Ecol*
1124 *Appl* 14:S127–S138.
- 1125 Neckel-Oliveira S, Gascon C (2006) Abundance, body size and movement patterns of a
1126 tropical tree frog in continuous and fragmented forests of the Brazilian Amazon. *Biol*
1127 *Conserv* 128:308–315.
- 1128 Nessimian JL, Venticinque EM, Zuanon J, De Marco P, Gordo M, Fidelis L, Batista J, Juen L
1129 (2008) Land use, habitat integrity, and aquatic insect assemblages in central
1130 Amazonian streams. *Hydrobiol* 614:117–131.
- 1131 Nogueira EM, Fearnside PM, Nelson BW, Barbosa RI, Keizer EWH (2008) Estimates of
1132 forest biomass in the Brazilian Amazon: New allometric equations and adjustments to
1133 biomass from wood-volume inventories. *For Ecol Manage* 256:1853-1857.
- 1134 Nogueira EM, Fearnside PM, Nelson BW, França MB (2007) Wood density in forests of
1135 Brazil's 'arc of deforestation': Implications for biomass and flux of carbon from land-
1136 use change in Amazonia. *For Ecol Manage* 248:119-135.
- 1137 Nogueira EM, Nelson BW, Fearnside PM (2005) Wood density in dense forest in central
1138 Amazonia, Brazil. *For Ecol Manage* 208:261-286.
- 1139 Norden N, Mesquita R, Bentos T, Chazdon R, Williamson GB (2011) Contrasting community
1140 compensatory trends in alternative successional pathways in central Amazonia. *Oikos*

- 1141 120:143-151
- 1142 Oliveira de AA, Mori SA (1999) A central Amazonian terra firme forest. I. High tree species
1143 richness on poor soils. *Biodiv Conserv* 8:1219–1244.
- 1144 Peres CA (2005) Why we need megareserves in Amazonia. *Conserv Biol* 19, 728–733.
- 1145 Peres CA, Gardner TA, Barlow J, Zuanon J, Michalski F, Lees A, Vieira I, Moreira F, Feeley
1146 KJ (2010) Biodiversity conservation in human-modified Amazonian forest landscapes.
1147 *Biol Conserv* 143:2314–2327.
- 1148 Phillips OL, Malhi Y, Higuchi N, Laurance WF, Nunez PV, Vasquez RM, et al. (1998)
1149 Changes in the carbon balance of tropical forests: Evidence from long-term plots.
1150 *Science* 282:439-442.
- 1151 Pimm SL (1998) The forest fragment classic. *Nature* 393:23–24.
- 1152 Pimm SL, Jenkins C (2005) Sustaining the variety of life. *Sci Amer* September:66–73.
- 1153 Powell AH, Powell GVN (1987) Population dynamics of male euglossine bees in Amazonian
1154 forest fragments. *Biotropica* 19:176–179.
- 1155 Puerta R (2002) Regeneração arbórea em pastagens abandonadas na região de Manaus em
1156 função da distância da floresta contínua. *Scientia Forestalis* 62:32–39.
- 1157 Quintela CE (1985) Forest Fragmentation and Differential Use of Natural and Man-made
1158 Edges by Understory Birds in Central Amazonia. M.Sc. Thesis, University of Illinois,
1159 Chicago.
- 1160 Quintero I, Roslin T (2005) Rapid recovery of dung beetle communities following habitat
1161 fragmentation in Central Amazonia. *Ecology* 86:3303–3311.
- 1162 Radtke MG, da Fonseca C, Williamson GB (2008) Forest fragment size effects on dung
1163 beetle communities. *Biol Conserv* 141:613–614.
- 1164 Rego F, Venticinque EM, Brescovit A (2007) Effects of forest fragmentation on four *Ctenus*
1165 spider populations (Araneae, Ctenidae) in central Amazonia, Brazil. *Stud Neotrop*
1166 *Fauna Environ* 42:137–144.
- 1167 Ribeiro MBN, Bruna EM, Mantovani W (2009) Influence of post-clearing treatment on the
1168 recovery of herbaceous plant communities in Amazonian secondary forests. *Restor*
1169 *Ecol* 18:50–58.
- 1170 Rocha R, López-Baucells A, Groenenberg M, Silva I, Farneda F, Bobrowiec P, Palmeirim
1171 JM, Meyer CFJ (2013) Effects of forest fragmentation on central Amazonian bats.
1172 Paper presented at the 16th International Bat Research Conference & 43rd North
1173 American Symposium on Bat Research, San José, Costa Rica.
- 1174 Salazar LF, Nobre CA, Oyama MD (2007) Climate change consequences on the biome
1175 distribution in South America. *Geophys Res Lett* 34:L09708.
- 1176 Sampaio EM, 2000. Effects of Forest Fragmentation on the Diversity and Abundance Patterns
1177 of Central Amazonian Bats. Ph.D. Dissertation, University of Tübingen, Germany.
- 1178 Sampaio EM, Kalko E, Bernard E, Rodriguez-Herrera B, Handley C, (2003) A biodiversity
1179 assessment of bats (Chiroptera) in a tropical lowland forest of central Amazonia,
1180 including methodological and conservation considerations. *Stud Neotrop Fauna*
1181 *Environ* 28:17–31.
- 1182 Santos BA, Arroyo-Rodríguez V, Moreno CE, Tabarelli M (2010) Edge-related loss of tree
1183 phylogenetic diversity in the severely fragmented Brazilian Atlantic forest. *PLoS ONE*
1184 5:e12625.
- 1185 Scariot A (1999) Forest fragmentation effects on diversity of the palm community in central
1186 Amazonia. *J Ecol* 87:66–76.
- 1187 Scariot A (2001) Weedy and secondary palm species in central Amazonian forest fragments.
1188 *Rev Bot Brasil* 15:271–280.
- 1189 Sizer N, Tanner EVJ (1999) Responses of woody plant seedlings to edge formation in a
1190 lowland tropical rainforest. Amazonia. *Biol Conserv* 91:135–142.

- Sizer N, Tanner EVJ, Kossman-Ferraz I (2000) Edge effects on litterfall mass and nutrient concentrations in forest fragments in central Amazonia. *J Trop Ecol* 16:853–863.
- Skole DS, Tucker CJ (1993) Tropical deforestation and habitat fragmentation in the Amazon: satellite data from 1978 to 1988. *Science* 260:1905–1910.
- Sodhi NS, Koh LP, Brook BW, Ng P (2004) Southeast Asian biodiversity: an impending disaster. *Trends Ecol Evol* 19:654–660.
- Stickler CM et al. (2009) The potential ecological costs and cobenefits of REDD: A critical review and case study from the Amazon region. *Glob Change Biol* 15:2803–2824.
- Stouffer PC, Bierregaard RO (1995a) Effects of forest fragmentation on understory hummingbirds in Amazonian Brazil. *Conserv Biol* 9:1085–1094.
- Stouffer PC, Bierregaard RO (1995b) Use of Amazonian forest fragments by understory insectivorous birds. *Ecology* 76:2429–2445.
- Stouffer PC, Bierregaard RO, Strong C, Lovejoy TE (2006) Long-term landscape change and bird abundance in Amazonian rainforest fragments. *Conserv Biol* 20:1212–1223.
- Stouffer PC, Strong C, Naka LN (2008) Twenty years of understory bird extinctions from Amazonian rain forest fragments: consistent trends and landscape-mediated dynamics. *Divers Distrib* 15:88–97.
- Stratford JA, Stouffer PC (1999) Local extinctions of terrestrial insectivorous birds in Amazonian forest fragments. *Conserv Biol* 13:1416–1423.
- Terborgh J, Lopez L, Nuñez VP, Rao M, Shahabuddin G, Orihuela G, Riveros M, Ascanio R, Adler G, Lambert T, Balbas L (2001) Ecological meltdown in predator-free forest fragments. *Science* 294:1923–1926.
- Ter Steege H, Pitman NCA, Sabatier D, Baraloto C, Salomao RP, Ernesto Guevara J, Phillips OL et al. (2013) Hyperdominance in the Amazonia flora. *Science* 342:326–334.
- Timo TPC (2003) Influência de fragmentação e matriz sobre a comunidade de mamíferos de médio e grande porte em uma floresta de terra firme na Amazônia central. M.Sc. Thesis, National Institute for Amazonian Research (INPA), Manaus, Brazil.
- Tocher M, Gascon C, Zimmerman BL (1997) Fragmentation effects on a central Amazonian frog community: a ten-year study. In: Laurance WF, Bierregaard RO (Eds), *Tropical forest remnants: Ecology, management, and conservation of fragmented communities*. University of Chicago Press, Chicago, pp 124–137.
- Trancoso R (2008) Hydrological Impacts of Deforestation in Small Catchments in Brazilian Amazonia. M.Sc. Thesis, National Institute for Amazonian Research (INPA), Manaus, Brazil.
- Uriarte M, Bruna EM, Rubim P, Anciães M, Jonckheere I (2010) Effects of forest fragmentation on the seedling recruitment of a tropical herb: assessing seed vs. safe-site limitation. *Ecology* 91:1317–1328.
- Van Houtan KS, Pimm SL, Bierregaard RO, Lovejoy TE, Stouffer PC (2006) Local extinctions in Amazonian forest fragments. *Evol Ecol Res* 8:129–148.
- Van Houtan KS, Pimm SL, Halley J, Bierregaard RO, Lovejoy TE (2007) Dispersal of Amazonian birds in continuous and fragmented forest. *Ecol Lett* 10:219–229.
- Vasconcelos HL, Luizão FJ (2004) Litter production and litter nutrient concentrations in a fragmented Amazonian landscape: edge and soil effects. *Ecol Applic* 14:884–892.
- Wilcox BA, Murphy DD (1985) Conservation strategy: the effects of fragmentation on extinction. *Am Nat* 125:879–887.
- Wieland LM, Mesquita RC, Bobrowiec PED, Bentos TV, Williamson GB (2011) Seed rain and advance regeneration in secondary succession in the Brazilian Amazon. *Trop Conserv Sci* 4:300–316.
- Williamson GB, Laurance WF, Oliveira A, Delamonica P, Gascon C, Lovejoy TE, Pohl L (2000) Amazonian wet forest resistance to the 1997–98 El Niño drought. *Conserv Biol*

1241 14:1538–1542.

1242 Williamson GB, Longworth JB, Bentos TV, Mesquita RC (2014) Convergence and
1243 divergence in alternative successional pathways in Central Amazonia. *Plant Ecol*
1244 *Diversity* 7:341–348.

1245 Williamson GB, Mesquita R (2001) Effects of fire on rain forest regeneration in the Amazon
1246 Basin. In: Laurance WF, Bierregaard RO (Eds), *Tropical forest remnants: Ecology,*
1247 *management, and conservation of fragmented communities.* University of Chicago
1248 Press, Chicago, pp 325–334.

1249 Zartman CE (2003) Forest fragmentation effects on epiphyllous bryophyte communities in
1250 central Amazonia. *Ecology* 84:948–954.

1251 Zartman CE, Nascimento HEM (2006) Are patch-tracking metacommunities dispersal
1252 limited? Inferences from abundance-occupancy patterns of epiphylls in Amazonian
1253 forest fragments. *Biol Conserv* 127:46–54.

1254 Zartman CE, Shaw AJ (2006) Metapopulation extinction thresholds in rainforest remnants.
1255 *Am Nat* 167:177–189.

1256
1257
1258

FIGURE CAPTIONS

Figure 1. Map of the Biological Dynamics of Forest Fragments Project in central Amazonia.

Figure 2. The diversity of edge-effect phenomena studied at the BDFFP and the distance to which each was found to penetrate into fragment interiors (after Laurance et al. 2002).

Figure 3. Forest plots affected by two or more nearby edges (plot center <100 m from edge) suffer greater tree mortality (A) and have a higher density (B) and species richness (C) of disturbance-loving pioneer trees than do plots with just one nearby edge. Values shown are the mean \pm SD (after Laurance et al. 2006a).

Figure 4. Increasing divergence of tree-community composition in three fragmented Amazonian landscapes. Tree communities in forest-edge plots (<100 m from the nearest edge) are shown before forest fragmentation and 13–18 years after fragmentation, based on a single ordination of all plots and censuses in the study area. The ordination used importance values for all 267 tree genera found in the plots (after Laurance et al. 2007).

Fig. 5. Tree genera that increase in forest fragments (those with positive values on the X axis) tend to have lower wood density than do those that decline in fragments (those with negative values on the X axis)(after Laurance et al. 2006b).

Fig. 6. Striking increases in the density of 52 species in nine pioneer or early successional genera (*Annona*, *Bellucia*, *Cecropia*, *Croton*, *Goupia*, *Jacaranda*, *Miconia*, *Pourouma*, *Vismia*) near forest edges in the BDFFP study area (after Laurance et al. 2006a).