

This file has been cleaned of potential threats.

If you confirm that the file is coming from a trusted source, you can send the following SHA-256 hash value to your admin for the original file.

b4664001ebe5e172e568d055f9bdf8e8a75318c1b7f2d96e09cd9496f7fd91e

To view the reconstructed contents, please SCROLL DOWN to next page.

**The text that follows is a PREPRINT.  
O texto que segue é um PREPRINT.**

**Please cite as:**

**Favor citar como:**

Ferrante, L., D. Rojas-Ahumada, M.  
Menem & P.M. Fearnside. 2023.  
**Climate change in the  
Central Amazon and its  
impacts on frog populations.**  
*Environmental Monitoring and  
Assessment* (in press).

ISSN: Electronic 1573-2959; Print 0167-6369

Copyright: Springer

The original publication will be available at  
O trabalho original estará disponível em:

<https://www.springer.com/journal/10661>

# Climate change in the central Amazon and its impacts on frog populations

Lucas Ferrante<sup>1, 2,\*</sup>, Diana Rojas-Ahumada<sup>2</sup>, Marcelo Menin<sup>1,†</sup> & Philip Martin Fearnside<sup>3</sup>

<sup>1</sup>Universidade Federal do Amazonas (UFAM), Manaus, Amazonas, Brazil.

<sup>2</sup>Programa de Pós-Graduação em Ecologia, Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Amazonas, Brazil.

<sup>3</sup>Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Amazonas, Brazil.

<sup>†</sup>Posthumous author.

\*Correspondence to: [lucasferrante@hotmail.com](mailto:lucasferrante@hotmail.com)

Lucas Ferrante: [lucasferrante@hotmail.com](mailto:lucasferrante@hotmail.com)

Diana Rojas-Ahumada: [draplucas@gmail.com](mailto:draplucas@gmail.com)

Philip Martin Fearnside: [pmfearn@inpa.gov.br](mailto:pmfearn@inpa.gov.br)

**Abstract.** Frog population declines have already been observed in the central Amazon even for common species that are considered not to be in danger of extinction. The Amazon is close to its limit of tolerated deforestation, and parts of the forest have already been modified by climate change, which raises questions about how the fauna in these areas would adapt to climate changes by the middle and the end of this century. In this study we used population density data on seven species of Amazonian frogs and analyzed the relationship between the activity of these species and temperature, precipitation, and relative humidity. We also used the least-squares method with logarithmic models to assess whether climate change projected by the Intergovernmental Panel on Climate Change (IPCC) would be an indicator of the population dynamics of these species. Our results suggest that even common species may be may experience population declines and extinction in the next decades due to climate changes.

**Key Words:** Anurans; Amphibian decline; Amazon Forest; biodiversity crises; bioindicators; Climate change; deforestation; Tropical Forest; Umbrella species.

Number of the words: Text: 4228; Total: 7274

## 42 INTRODUCTION

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

In Amazonia, as in other parts of the world, the fate of amphibians serves as a harbinger of loss of biodiversity and much wider environmental destruction (Zipkin et al. 2020; Strona and Bradshaw 2022). Climate change has altered the geographical distributions of multiple anuran species around the world (e.g., Vieira et al. 2018; Hu et al. 2019). Climatic disturbance events have been responsible for the sudden population decline and extinction of several anuran species in Central America (Pounds and Crump 1994; Pounds et al. 1999). Other factors acting in synergy with climatic events have caused local extinctions in the Atlantic Forest of Brazil (Heyer et al. 1988; Weygoldt 1989; Eterovich et al. 2005; Ferrante et al. 2019).

In the central Amazon some anuran species would be unable to migrate to climatically suitable locations, as in the case of the endemic central-Amazonian frog *Atelopus manauensis* (Jorge et al. 2020). This is especially so since Amazonian rivers are geographical barriers for various vertebrate groups, such as birds (Ribas et al. 2012; Ferreira et al. 2016; Braga et al. 2022) and also amphibians and squamates (Moraes et al. 2016). Barriers can also be imposed by land-use change, particularly the growing areas of deforestation that leave remaining areas of natural forest as islands surrounded by a landscape that is hostile to the forest's endemic frogs (Ferrante et al. 2017, 2019), and even to some species with significant dispersal and colonization ability (Ferrante et al. 2020).

Deforestation is expanding from Manaus into surrounding areas of the central Amazon (Ramos et al. 2018; Santos et al. 2022), and this transformation would be greatly accelerated by the planned reconstruction of Highway BR-319 connecting Manaus to Brazil's "arc of deforestation" - the highly deforested strip of land along the southern and eastern edges of Brazil's Amazon region (Ferrante et al. 2021a). Other processes creating barriers include forest degradation through fire and consequent expansion of savannas (Sales et al. 2020; Flores and Holmgren 2021). When climate change occurs in areas isolated by barriers, the species must either adapt to local environmental changes or become extinct (Quental and Marshall 2013; Ferrante et al. 2023) unless they succeed in colonizing new ecologically suitable habitats. Climate change is considered to be one of the greatest threats to amphibians (Stuart et al. 2008; Bishop et al. 2012), which is the most threatened group among vertebrates (Baillie et al. 2010; Bishop et al. 2012). Characteristics that make frogs vulnerable to climate change include the fact that their skin is permeable, and they have exposed eggs and embryos - most species have a free-swimming larval stage in the life cycle (Blaustein et al. 1994, 2001; Blaustein and Kiesecker 2002; Blaustein and Bancroft 2007; Duellman and Trueb 1996; Stebbins and Cohen 1995).

Climate change has already affected different taxonomic groups in the central Amazon, especially in valley and stream areas, as shown by more than 20 years of monitoring (Costa et al. 2020). For some locations in Brazil, anthropogenic climate change in synergy with landscape change have been shown to threaten anuran populations and cause local extinctions (Ferrante et al. 2019) even more than invasive species or lethal pathogens (Ferrante et al. 2020). This means that anthropogenic climate change should be a priority for studies of threats to Brazilian amphibians (Verdade et al. 2012). In addition, structural changes in the landscape (such as deforestation, forest fragmentation, edge effects and neighboring agricultural crops) can alter the microclimate of amphibian habitats, leading to subtle increases in temperature and losses of moisture that can affect population density, species richness and community composition (Urbina-Cardona 2006; Ferrante et al. 2017).

92 Amphibians are extremely vulnerable to climate change in the Amazon  
93 (Vasconcelos et al. 2018), and the distribution of anuran taxa is directly influenced by  
94 the rainy season and by proximity to humid environments, such as valleys and streams  
95 (Moraes et al. 2016), which are already altered by ongoing climate change (Costa et al.  
96 2020). The present study aims to show how Amazonian frog species are responding to  
97 climate change. We hypothesize that ongoing climate change could even threaten the  
98 populations of common species that are now abundant and have so far not been  
99 considered to be threatened.


## 100 **METHODS:**

### 101 *Study Site*

102  
103  
104  
105 The region denominated here as the “central Amazon” (Fig. 1) is bounded to the  
106 east by the border between the states of Amazonas and Pará (a NE-SW diagonal from  
107 approximately 57 to 60° W longitude), thus avoiding the “dry corridor” that crosses the  
108 Amazon River at Santarém, Pará. To the west, the central Amazon can be considered  
109 bounded at 66° W Longitude, thus avoiding the area with a dry season < 2 months in  
110 length in the NW corner of Brazilian Amazonia. To the north it is bounded by the  
111 equator, and to the south at approximately 6° S latitude, thus avoiding the Humaitá  
112 savanna and areas with a dry season > 3 months in length. This region encompasses a  
113 range of ecoregions (Olson et al. 2001; Fearnside 2023), with a predominance of dense  
114 ombrophilous forest and is in the morphoclimatic domain classified as “equatorial  
115 forested lowlands,” with average temperatures of 24 to 27°C (Ab'Sáber 2003). The  
116 central Amazon is vulnerable to climate change and is becoming increasingly  
117 susceptible to forest fires, especially in El Niño years (Reis et al. 2021).

118 Data used in the present study were obtained in the Adolpho Ducke Forest  
119 Reserve, in the municipality of Manaus in the central Amazon. The Adolpho Ducke  
120 Forest Reserve is located in *terra firme* (upland) forest. Although these forests are not  
121 seasonally flooded by large rivers (Braga 1979), they have many permanent and  
122 temporary pools that are used by amphibians for reproduction (Menin et al. 2008; Najar  
123 and Ferrante 2018). The study area has an average annual temperature of 26 °C and  
124 annual rainfall ranging from 1750 to 2500 mm (Oliveira et al. 2008) - a sampling  
125 interval that encompasses the rainfall regimes of the different ecoregions included in the  
126 central Amazon (Ab'Sáber 2003). All of the landscape studied here has the same forest  
127 type (dense ombrophilous forest) and is the best-studied area in the central Amazon.  
128 The Adolpho Ducke Forest Reserve has a large sampling effort in terms of biodiversity  
129 and physical and climatic variables and, as a permanently protected area, the sampled  
130 sites are not susceptible to impacts other than anthropogenic climatic effects (Oliveira et  
131 al. 2008; Magnusson et al. 2013).

132



**Figure 1.** Map of ecoregions in the central Amazon and Adolpho Ducke Forest Reserve (ADFR).

### **Sampling design**

Declines in anuran populations can be caused by the lengthening of the dry season and by reduced rainfall in the wet season; these changes lower the activity of individuals, and a substantial drop in a single season can affect the recruitment of new individuals in the population and decrease the population density in subsequent years (Ferrante et al. 2023). Population data for the seven species in the present study were collected in five samplings: November-December 2002; March-May 2003; November-December 2003; February to May 2004 (Data from Menin et al. 2008). The sampling periods included both the dry season, from June to November, and the rainy season, from December to May (Ribeiro 1976), thereby allowing us to associate fluctuations in the active frog populations with climatic fluctuations in the central Amazon.

This means that alterations in the dry and wet seasons caused by global climate change can shape anuran population dynamics in the long term, since there is not enough time for adaptation, as has already been observed for frogs in Brazil's Atlantic Forest (Ferrante et al. 2023). In the present study we use data from monitoring anuran populations over a period that covers both the dry and wet seasons and that can capture population oscillations due to local climatic variations. Knowing the temporal dynamics of the species and the abundance variation in the observed samples, it is possible to assess the degree of dispersion of the probability values. The range of variation therefore allows testing a climate-change scenario and obtaining a model that allows projection of how the species will behave within the limits of this model.

Population census data were analyzed for seven anuran species (*Atelopus manauensis* (n=21), *Leptodactylus pentadactylus* (n=63), *Leptodactylus rhodomystax* (n=15), *Osteocephalus oophagus* (n=3222), *Pristimantis fenestratus* (n=6702), *Synapturanus mirandariberoi* (n=1459) and *Synapturanus salseri* (n=996)) and average

162 climatological data (average temperature, maximum temperature, relative humidity of  
163 the air and precipitation) for the different collection periods. Linear and logarithmic  
164 regressions were used to test the relationship between the population activity of these  
165 seven species in relation to climatic oscillations.

166 Sampling was performed by visual and auditory means simultaneously (Crump  
167 and Scott 1994) considering only mature individuals. The measure of “population  
168 density” is the abundance of individuals per sampling period. Each sampling period  
169 lasted an average of 49 days with two observers searching the plots for 2 hours per day,  
170 each sampling period having 196 hours of observation (49 days  $\times$  2 hours per day  $\times$  2  
171 observers). The plots were 250  $\times$  40 m (1 ha) in area (Magnusson et al. 2005), and in the  
172 shorter sampling periods the number of plots was increased to keep the sampling effort  
173 constant. Due to gaps in the climatic data and the aggregation of the anuran population  
174 data by season in Menin et al. (2008), we used the values for temperature and relative  
175 humidity of the air provided by the National Institute of Meteorology (INMET 2023);  
176 precipitation data were collected during the anuran collection itself (See. Appendix S1).

177 Both analyses of continuous data and of aggregated data have been used in  
178 ecological studies, with losses and gains between these two types of analysis depending  
179 on the question to be answered (Gotelli and Ellison 2004; Magurran 2004; Magnusson  
180 and Mourão 2005). Aggregated individual abundance data are common in ecological  
181 analyses (Gotelli and Ellison 2004). Here, the aggregation of biological data by season  
182 makes it possible to visualize the population response of each species to climate change  
183 in each season. Climatic anomalies that may occur on certain days, or local stochastic  
184 effects in certain plots, do not tend to generate outliers or sampling artifacts in the data  
185 analysis. This also allows us to have greater reliability in population density projections  
186 in relation to future climate change (Mills 2013) because the data are for censuses of  
187 active individuals over a long period with an established climate pattern. Grouping the  
188 data avoids population fluctuations based on the gradual change in temperature or  
189 humidity through the course of the day.

## 190 *Statistical analyses*

191 We performed simple linear regressions, together with tests of homoscedasticity  
192 using the Bartlett test (Bartlett 1937) and Shapiro Wilk normality tests (Shapiro and  
193 Wilk 1965) (See. Appendix S2 and S3). Subsequently, data on the abundance of  
194 individuals in relation to climate variables were projected through logarithmic  
195 regressions using the least-squares method (Luenberger 1997; Tang and Wang 2001) in  
196 relation to the annual progression of climate variables predicted for this area from 2002  
197 to 2100 by the RCP 8.5 scenario of the IPCC’s fifth assessment report (AR5) (IPCC  
198 2014; 2022; Magrin et al. 2014). We used a natural logarithm distribution. Least squares  
199 can be derived as a method-of-moments estimator (Luenberger 1997) and are used here  
200 to estimate the moments of the frog population declines assuming the progression of  
201 climate change projected under the RCP 8.5 scenario (IPCC 2014; Magrin et al. 2014).  
202 We only tested the average and maximum effect of temperature on the species since an  
203 increase in temperature is expected by the RCP 8.5 model. Annual estimates under the  
204 RCP 8.5 scenario were taken from Science on a Sphere (2023). These models are well-  
205 suited for calculating local changes based on the least-squares method (Tang and Wang  
206 2001). This method has advantages over other regression tests commonly used in  
207 ecology due to its capacity to generate scenarios that enlarge the range of projections  
208 (Carrascal et al. 2009). All analyses were run in Statistica 8 software (Statsoft 2007).

210


211 **RESULTS:**

212

213 Linear regressions (Fig. 2) showed that at least one of the climatic variables  
214 significantly influenced the activities of three of the seven species studied: *Atelopus*  
215 *manauensis* in relation to the average temperature of the air (n=21, p = 0.01, R<sup>2</sup> = 0.87);  
216 *Leptodactylus rhodomystax* in relation to the average temperature of the air (n=15, p =  
217 0.01, R<sup>2</sup> = 0.91) and; *Synapturanus mirandaribeiroi* in relation to the precipitation  
218 (n=1459, p = 0.05, R<sup>2</sup> = 0.77).

219 Results for two additional species are suggestive of climate effects but were not  
220 significant at the p<0.05 level: *Leptodactylus pentadactylus* with a value of activity of  
221 this species with a suggestive relation to precipitation (n=996, p = 0.07, R<sup>2</sup> = 0.70). In  
222 addition, *Synapturanus mirandaribeiroi* had a suggestive relation to precipitation  
223 average temperature of the air (n=1459, p = 0.07, R<sup>2</sup> = 0.69), in addition to its  
224 significant relation to precipitation.






225  
226  
227  
228  
229  
230  
231

**Figure 2.** Relationships between the population density (calling individuals per sampling period) of the seven target species in the study and the climate variables for the collection area.

Due to lack of data on the thermal tolerance thresholds of species that responded positively to temperature increase, we did not project population trends for these species. Logarithmic regressions indicated that *L. rhodomystax* would have reduced

232 activity under the climate projected for the middle and the end of the current century  
 233 based on the relationships between population density and climatic variables (Fig. 3).  
 234 An increase in the annual mean temperature in the frogs' habitat to a level above 30 °C  
 235 tends to eliminate the activity of *L. rhodomystax*. Increases of 2 to 4 °C in the annual  
 236 average temperature in the central Amazon would decrease the activity of this species,  
 237 causing lower recruitment of individuals and population declines.  
 238  
 239  
 240



241  
 242 **Figure 3.** Effect of projected climate change on *L. rhodomystax* population densities per hectare in  
 243 response to increases in the annual average temperature in the central Amazon. The maps show the  
 244 projected climate in the mid-21<sup>st</sup> century (2050) and the late 21<sup>st</sup> century (2100) (Magrin et al. 2014).  
 245

246 Our results point to a decrease in the density (calling individuals per hectare) of  
 247 *L. rhodomystax* due to increased average annual temperature, with the possibility of  
 248 extinction of the species in the next 20 years. These results indicate that the density of  
 249 active individuals would decrease substantially in the next decades. Monthly  
 250 fluctuations in humidity and temperature naturally occur in the central Amazon, causing  
 251 the density of active individuals indicated here to vary over the course of the year. The  
 252 reductions indicated here are per hectare at the location of the plots, and declines in  
 253 other populations are likely because of reductions in the distribution areas of these  
 254 species.  
 255

## 256 DISCUSSION:

257  
 258 Anurans in the central Amazon had a strong response to climate and can be  
 259 considered as bioindicators of climate change. Our results indicate that species with  
 260 different ecological habits and adaptations may respond differently to certain climatic  
 261 variables (see methods), and their activity may be negatively affected by both the  
 262 average and the maximum temperatures, since changes in climate range affect tolerance  
 263 limits of frogs, as is the case for the observed climatic oscillations in the central Amazon  
 264 at present. The different thermal sensitivities of anuran species point to the need to  
 265 classify species according to their ecological habits in studies at the landscape level, as  
 266 in the studies by Urbina-Cardona et al. (2006) and Ferrante et al. (2017).  
 267

268 We note that the RCP 8.5 scenario from the IPCC's fifth assessment report  
 269 (AR5) (IPCC 2014) has been criticized as being overly pessimistic in its emissions  
 assumptions for the remainder of this century, especially with regard to use of coal (e.g.,

270 Ritchie and Dowlatabadi 2017). However, the RCP 8.5 scenario's best estimate for  
271 global mean temperature increase by 2100 (4.8 °C relative to the 1850–1900 mean) is  
272 only slightly higher than the SSP5-8.5 estimate in the AR6 (4.4 °C) (IPCC 2022). While  
273 restraining global emissions to levels below those used in these scenarios is hoped to  
274 occur, analyses assuming unrestrained emissions have value in illustrating the  
275 consequences of continued insufficient action in mitigating global warming. While  
276 assumptions regarding the use of coal create upward bias, these scenarios also contain  
277 biases in the downward direction by not including a variety of “indirect” emissions  
278 sources (e.g., Barros and Fearnside 2019). In addition, with the increase in the use of  
279 fossil fuels (Ferrante and Fearnside 2023) and the success of “ruralist” politicians in  
280 Brazil in dismantling many of the protections of the Amazon forest (Ferrante and  
281 Fearnside 2019; Ferrante 2023), which doubled carbon emissions in the Amazon (Gatti  
282 et al. 2023), make the most pessimistic scenarios today, such as RCP 8.5, conservative  
283 for the climate change expected in the region.

284 Ongoing climate change has already led to an increase in annual mean  
285 temperature in the Amazon rainforest (Marengo et al. 2021), with forecasts of a  
286 continued increase reaching up to 8 °C in some areas of the Amazon by the end of the  
287 century (IPCC 2014, 2022). In addition to the expected changes in average temperature  
288 precipitation and wind patterns, extreme weather events are expected to increase greatly  
289 both in terms of frequency and intensity (IPCC 2014, 2022). According to our data,  
290 these climate changes may negatively affect the population dynamics of frogs in the  
291 central Amazon.

292 Data on population declines and local extinctions of frogs caused by climate  
293 change in the Amazon region are still scarce (Stuart et al. 2008), although some  
294 population declines have already been observed due to other threats, such as the use of  
295 herbicides (Ferrante and Fearnside 2020a). The data presented here shed light on how  
296 Amazonian frog populations may react in the face of climate change by the end of this  
297 century.

298 Physiological studies have shown that some Brazilian amphibians lack adaptive  
299 plasticity even to seemingly small increases in temperature, negatively affecting their  
300 physiological performance in the larval stage (Longhini et al. 2021). This reinforces the  
301 hypothesis of declines and extinctions in the face of climate change. The decline of  
302 amphibian populations is also related to climate change due to ultraviolet radiation  
303 (Blaustein and Kiesecker 2002; Blaustein et al. 1994, 2001). This could prove to be  
304 catastrophic for local anuran populations because higher incidence of ultraviolet  
305 radiation is expected for the central Amazon (IPCC 2022). Amphibian eggs are sensitive  
306 to increases in ultraviolet radiation, consequently causing high mortality of embryos  
307 (Blaustein et al. 1994). This would have a profound impact, especially on diurnal  
308 species such as *A. manauensis* (Menin et al. 2008). The climatic optimum for this  
309 species is currently in a range of warm average temperatures (Fig. 2), but the behavior of  
310 the species may change given the high temperatures predicted in climate-change  
311 scenarios for the central Amazon.

312 Extreme weather events have already caused the extinction of several amphibian  
313 species in Central and South America (Pounds and Crump 1994; Pounds et al. 1999;  
314 Stuart et al. 2008). The frequency of extreme drought events (due to strong El Niño  
315 episodes) has increased in the Amazon Basin since the 1970s, departing from the long-  
316 term pattern that had predominated since 1901 (Paredes-Trejo et al. 2021).  
317 “Unprecedented” droughts are projected to occur in Amazonia in the coming decades  
318 (Kay et al. 2022). Climate change that is already underway across South America,  
319 including tropical forests such as the Amazon and Atlantic Forests (Stuart et al. 2008),

320 motivated the inclusion of climate change in the Brazilian Amphibian Conservation  
321 Action Plan (Verdade et al. 2012).

322 Future scenarios proposed through modeling portend a worldwide loss of  
323 amphibian species through climate change, with some groups being more threatened  
324 than others (Loyola et al. 2013). Our results suggest that even common species, such as  
325 *L. rhodomystax*, can be expected to undergo reductions in their population densities as a  
326 result of the increase of temperature (by 2100 under the RCP 4.5 scenario, which, unlike  
327 the RCP 8.5 scenario, assumes substantial reductions in global greenhouse-gas  
328 emissions). We should therefore expect decreases in the activity levels of these anurans  
329 in the central Amazon. This would result in lower recruitment of individuals and  
330 population declines of these species.

331 The impact of increases in the annual mean values of climatic variables are  
332 much more serious than only anuran population declines, as climate change affects  
333 much more than the activity of frogs (Costa et al. 2020). There are also threats to the  
334 central Amazon from burning (Fearnside 2021), land grabbing (Ferrante et al. 2021b),  
335 illegal logging (Andrade et al. 2021), illegal mining (Ferrante and Fearnside 2022a), and  
336 the expansion of agriculture and cattle ranching (Ferrante and Fearnside 2018, 2020a,  
337 2020b, 2022b; Ferrante et al. 2021c). These multiple threats have substantial impacts on  
338 forest structure and consequently affect the microclimate, which in turn affects both the  
339 richness and the abundance of amphibians (Urbina-Cardona et al. 2006; Ferrante et al.  
340 2017). Shifts in the amplitude of variation in climatic variables may force species  
341 generally to the limits of their physiological tolerance and restrict their geographical  
342 distributions (Mills 2013). Loss of certain species in an ecosystem can collapse trophic  
343 chains and unbalance the dynamics of populations of other species, causing extinctions  
344 at the local and regional levels or for the species as a whole (Zipkin et al. 2020; Strona  
345 and Bradshaw 2022). Endemic species, such as *A. manauensis*, are likely to be the most  
346 affected due to their restricted geographical distributions (Jorge et al. 2020). The species  
347 in the present study can be considered to constitute an umbrella species group whose  
348 conservation would confer protection to a large number of naturally co-occurring  
349 species. The loss of amphibian species around the world has triggered a cascade effect,  
350 which has affected other taxonomic groups, mainly predators, as is already seen in  
351 Neotropical snakes (Zipkin et al. 2020). Our results warn of a potential widespread loss  
352 of biodiversity in the central Amazon by 2050 and even greater impacts by 2100, as  
353 anurans are bioindicators for declines in other taxonomic groups (Zipkin et al. 2020). It  
354 is crucial that Brazil adopt measures to mitigate climate change and to protect  
355 biodiversity.

356 The threat to Amazon biodiversity (by climate change documented here) is  
357 embedded in a context of widespread destruction of the region's native ecosystems by  
358 deforestation, logging, forest fires, mining, dams, and other economic ventures (e.g.,  
359 Fearnside 2021). These processes accelerated under the 2019-2022 Jair Bolsonaro  
360 presidential administration in Brazil, with the dismantling of the country's  
361 environmental agencies, including hundreds of changes in internal operating rules in  
362 ways that impede enforcement of environmental regulations, multiple legislative  
363 changes loosening environmental restrictions, and a constant discourse denying  
364 scientific results on climate, deforestation, and fire and suggesting that environmental  
365 crimes would be ignored or pardoned (Ferrante and Fearnside 2019; Diele-Viegas et al.  
366 2021; da Silva and Fearnside 2022). Many of these setbacks will have lasting effects  
367 irrespective of current and future efforts to reverse them. Building roads, for example,  
368 allows migrations and unleashes processes that are largely outside of government  
369 control, a fact that is particularly relevant to the central Amazon given the advance of a

370 project to reconstruct Highway BR-319 connecting this relatively intact area to Brazil's  
371 "arc of deforestation" (Andrade et al. 2021; Ferrante et al. 2021a,b; Fearnside 2022).  
372 President Luiz Inácio Lula da Silva, who took office in January 2023, has promised to  
373 reduce deforestation, but restoring the control capacities of environmental agencies will  
374 not be enough because avoiding large-scale habitat loss in the central Amazon will also  
375 require blocking major projects such as the reconstruction of Highway BR-319 and  
376 opening oil and gas fields in the proposed Solimões Sedimentary Basin project (e.g.,  
377 Bustamante et al. 2023; Ferrante and Fearnside 2023; Vilani et al. 2023).

378 In addition to contributing to the climate change that would affect frogs and  
379 other groups, the deforestation provoked by BR-319 and its associated side roads would  
380 have severe impacts on biodiversity through habitat loss (Magnusson 2020; Ferrante et  
381 al. 2021a). Highway BR-319 would bring to the central Amazon the land-use change  
382 processes already present in Brazil's "arc of deforestation." This would be catastrophic  
383 for Amazonian amphibians. In addition to habitat loss and a contribution to the climate  
384 change that threatens these species, the highway is likely to act as a disperser of new  
385 pathogens that are lethal to amphibians, such as the fungus *Batrachochytrium*  
386 *dendrobatidis* (Bd) (Becker et al. 2016). Bd has already been recorded in the "arc of  
387 deforestation" (Becker et al. 2016), and Highway BR-319 would offer a direct route for  
388 the fungus to spread to the central Amazon. The massive population declines of  
389 amphibians for the central Amazon that the present study suggests because of projected  
390 climate change are therefore conservative due to the other simultaneous threats resulting  
391 from the BR-319 project. The BR-319 project's direct and indirect threats to frogs add  
392 to the many reasons indicating that the project should not be undertaken (Fearnside  
393 2022).

394 Control of Brazil's National Congress by the "ruralist" voting block has been  
395 increased by the 2022 elections (ClimaInfo 2022), suggesting likely passage in the  
396 coming months and years of a series of proposed laws further weakening environmental  
397 control, facilitating Amazonian land grabbing, and opening indigenous lands to  
398 agribusiness, mining and other activities by nonindigenous people (Ferrante and  
399 Fearnside 2021, 2022c; Ruaro et al. 2021, 2022). The combination of impacts from  
400 climate change and from habitat destruction by direct human action implies massive  
401 losses of Amazonian biodiversity (e.g., Joly et al. 2019). However, both Brazil's  
402 policies affecting deforestation and global accords on measures to contain climate  
403 change are subject to human decisions, and efforts must not be spared to avoid the bleak  
404 future that current trends imply.

405

## 406 **CONCLUSION:**

407

408 The population density of anurans in the central Amazon is strongly influenced  
409 by temperature, precipitation, and relative humidity. By the end of the 21<sup>st</sup> century,  
410 projected climate change would even affect the population dynamics of common species  
411 that today are considered to be out of danger, causing population declines and possibly  
412 local extinctions in many species through extreme climatic events. Habitat loss from  
413 deforestation and other direct anthropogenic impacts further increase the risks to frogs  
414 and other groups in the central Amazon.

415

## 416 **DECLARATIONS**

417

418 All authors have read, understood, and have complied as applicable with the statement  
419 on "Ethical responsibilities of Authors" as found in the Instructions for Authors and are

420 aware that, with minor exceptions, no changes can be made to authorship once the paper  
421 is submitted.

422

#### 423 **FUNDING DECLARATION**

424 L.F. thanks the National Council for Scientific and Technological Development  
425 (CNPq). This study was financed in part by the Coordenação de Aperfeiçoamento de  
426 Pessoal de Nível Superior (CAPES) – Finance Code 001. P.M.F. thanks CNPq (Proc.  
427 312450/2021-4) and the Brazilian Research Network on Climate Change (FINEP/Rede  
428 CLIMA, Proc. 01.13.0353-00).

429

#### 430 **CONFLICT OF INTEREST**

431 The authors declare no conflict of interest.

432

#### 433 **AUTHOR CONTRIBUTIONS**

434 L.F. designed the research; D.R. conducted fieldwork and examined material; L.F. conducted  
435 statistical analyses; L.F., D.R. and P.M.F. wrote the manuscript; L.F., D.R. and P.M.F. revised  
436 the manuscript.

437

#### 438 **DATA AVAILABILITY STATEMENT**

439 All data are available upon request to the corresponding author.

440

441 **Acknowledgments:** We are especially grateful to Professor Marcelo Menin of the  
442 Federal University of Amazonas, who kindly provided the data collected in the Adolpho  
443 Ducke Reserve and who was part of this study until he passed away from COVID-19 in  
444 2021; his contribution was invaluable to herpetology, and his death is an irreparable  
445 loss.

446

#### 447 **REFERENCES:**

448

449 Ab'Sáber A (2003) Os Domínios de Natureza no Brasil: Potencialidades Paisagísticas.  
450 Ateliê Editorial, São Paulo, Brazil. 160 pp.

451

452 Andrade M, Ferrante L, Fearnside PM (2021) Brazil's Highway BR-319 demonstrates a  
453 crucial lack of environmental governance in Amazonia. *Environmental Conservation*,  
454 48, 161-164. DOI: <https://doi.org/10.1017/S0376892921000084>

455

456 Baillie JEM, Griffiths J, Turvey ST, Loh J, Collen B (2010) Evolution lost: Status and  
457 trends of the world's vertebrates. Zoological Society of London, London, UK. 72 pp.

458

459 Bartlett MS (1937). Properties of sufficiency and statistical tests. *Proceedings of the Royal*  
460 *Statistical Society*, 160, 268–282.

461

462 Barros HS, Fearnside, PM (2019) Soil carbon is decreasing under “undisturbed”  
463 Amazonian forest. *Soil Science Society of American Journal*, 83, 1779-1785. DOI:  
464 <http://doi.org/10.2136/sssaj2018.12.0489>

465

466 Becker CG, Rodriguez D, Lambertini C, Toledo LF, Haddad CFB (2016) Historical  
467 Dynamics of *Batrachochytrium dendrobatidis* in Amazonia. *Ecography*, 39, 954-960.  
468 DOI : <https://onlinelibrary.wiley.com/doi/epdf/10.1111/ecog.02055>

469

- 470 Bishop PJ, Angulo A, Lewis JP, Moore RD, Rabb GB, Moreno JG (2012) The  
471 Amphibian Extinction Crisis - what will it take to put the action into the Amphibian  
472 Conservation Action Plan? S.A.P.I.EN.S., 5.2. URL:  
473 <http://journals.openedition.org/sapiens/1406>  
474
- 475 Blaustein AR, Bancroft BA (2007) Amphibian population declines: Evolutionary  
476 considerations. *BioScience*, 57, 437-444. DOI: <http://doi.org/10.1641/B570517>  
477
- 478 Blaustein AR, Kiesecker JM (2002) Complexity in conservation: Lessons from the  
479 global decline of amphibian populations. *Ecology Letters*, 5, 597-608. DOI:  
480 <https://doi.org/10.1046/j.1461-0248.2002.00352.x>  
481
- 482 Blaustein AR, Hoffman PD, Hokit DG, Kiesecker JM, Walls SC, Hays JB (1994) UV  
483 repair and resistance to solar UV-B in amphibian eggs: A link to population declines?  
484 *PNAS*, 1791-1795. DOI: <https://doi.org/10.1073/pnas.91.5.1791>  
485
- 486 Blaustein AR, Belden LK, Olson DH, Green DM, Root TL, Kiesecker JM (2001)  
487 Amphibian breeding and climate changes. *Conservation Biology*, 15, 1804-1809.  
488 <https://www.jstor.org/stable/3061281>  
489
- 490 Braga PIS (1979) Subdivisão fitogeográfica, tipos de vegetação, conservação e  
491 inventário florístico da floresta amazônica. *Acta Amazonica*, 9, 53-80. DOI:  
492 <https://doi.org/10.1590/1809-43921979094s053>  
493
- 494 Braga PLM, Borges SH, Peres CA, Loisele BA, Blake JG, Menger J, Bueno AS,  
495 Anciães M, Teófilo FH, Maximiano MFA, Souza AHN, Boss RL, Baccaro FB (2022)  
496 Connecting Amazonian historical biogeography and local assemblages of understory  
497 birds: Recurrent guild proportionality within areas of endemism. *Journal of*  
498 *Biogeography*, 49, 324-338. DOI: <https://doi.org/10.1111/jbi.14301>  
499
- 500 Bustamante MMC, Hipolito J, Delgado PGG, Ferrante L, Vale MM (2023) The future  
501 of Brazilian science. *Nature Human Behaviour*, 7, 825–827. DOI:  
502 <https://doi.org/10.1038/s41562-023-01597-7>  
503
- 504 Carrascal LM, Galván I, Gordo O (2009) Partial least squares regression as an  
505 alternative to current regression methods used in ecology. *Oikos*, 118, 681-690. DOI:  
506 <https://doi.org/10.1111/j.1600-0706.2008.16881.x>  
507
- 508 ClimaInfo (2022) Desmonte ambiental: Próximo Congresso será “mais boiadeiro” que o  
509 atual. ClimaInfo, 6 October 2022. <https://bit.ly/3rRCngm>  
510
- 511 Costa FRC, Zuanon JAS, Baccaro FB, Almeida JS, Menger JS, Souza JLP, Borba GC,  
512 Esteban EJ L, Bertin VM, Gerolamo CS, Nogueira A, Castilho CV (2020) Effects of  
513 climate change on central Amazonian Forests: A two decades synthesis of monitoring  
514 tropical biodiversity. *Oecologia Australis*, 24, 315-333. DOI:  
515 <https://doi.org/10.4257/oeco.2020.2402.07>  
516
- 517 Crump ML, Scott NJ (1994) Visual encounter surveys. Measuring and monitoring  
518 biological diversity. In: Standard Methods for Amphibians. Heyer, WR, Donnelly, MA,

- 519 McDiarmid RW et al. (eds). Smithsonian Institution Press, Washington, DC, USA. pp.  
520 84–92.  
521
- 522 da Silva MD, Fearnside, PM (2022) Brazil: Environment under attack. *Environmental*  
523 *Conservation*, 49(4), 203–205. DOI: <https://doi.org/10.1017/S0376892922000364>  
524
- 525 Diele-Viegas LM, Hipolito J, Ferrante L (2021) Scientific denialism threatens Brazil.  
526 *Science*, 374, 948-949. DOI: <https://doi.org/10.1126/science.abm9933>  
527
- 528 Duellman WE, Trueb L (1996) *Biology of Amphibians*. The Johns Hopkins University  
529 Press, Baltimore, MD, USA. 670 pp.  
530
- 531 Eterovick PC, Carnaval ACOQ, Borges-Nojosa DM, Silvano DL, Segalla MV, Sazima I  
532 (2005) Amphibian declines in Brazil: An overview. *Biotropica*, 37, 166-179.  
533 <https://www.jstor.org/stable/30043169>  
534
- 535 Fearnside PM (2021) Deforestation in Brazilian Amazonia. In: Wohl E (ed.) Oxford  
536 Bibliographies in Environmental Science. Oxford University Press, New York, NY,  
537 USA. DOI: <https://doi.org/10.1093/obo/9780199363445-0064>  
538
- 539 Fearnside PM (2022) Amazon environmental services: Why Brazil’s Highway BR-319  
540 is so damaging. *Ambio*, 51, 1367–1370. DOI: [https://doi.org/10.1007/s13280-022-](https://doi.org/10.1007/s13280-022-01718-y)  
541 [01718-y](https://doi.org/10.1007/s13280-022-01718-y)  
542
- 543 Fearnside PM (2023) South American natural ecosystems, status of. In: Reference  
544 Module in Life Sciences, Elsevier, Amsterdam, The Netherlands. DOI:  
545 <https://doi.org/10.1016/B978-0-12-822562-2.00134-1>  
546
- 547 Ferrante L (2023) Bills undermine Brazil’s environmental goals. *Science*, 381, 490-491.  
548 DOI: <https://doi.org/10.1126/science.adi9196>  
549
- 550 Ferrante L, Baccaro FB, Ferreira EB, Sampaio MFO, Santos T, Justino RC, Angulo A  
551 (2017) The matrix effect: how agricultural matrices shape forest fragment structure and  
552 amphibian composition. *Journal of Biogeography*, 44, 1911-1922. DOI:  
553 <https://doi.org/10.1111/jbi.12951>  
554
- 555 Ferrante L, Barbosa RI, Duczmal L, Fearnside PM (2021c) Brazil’s planned  
556 exploitation of Amazonian indigenous lands for commercial agriculture increases risk of  
557 new pandemics. *Regional Environmental Change*, 21, art. 81. DOI:  
558 <https://doi.org/10.1007/s10113-021-01819-6>  
559
- 560 Ferrante L, Andrade MBT, Leite L, Silva Junior CA, Lima M, Coelho Junior MG, Neto  
561 ECS, Campolina D, Carolino K, Diele-Viegas ML, Pereira EJAL, Fearnside PM  
562 (2021a) Brazil’s Highway BR-319: The road to the collapse of the Amazon and the  
563 violation of indigenous rights. *Die Erde*, 152, 65-70. DOI:  
564 <https://doi.org/10.12854/erde-2021-552>  
565
- 566 Ferrante L, Andrade MBT, Fearnside PM (2021b) Land grabbing on Brazil’s Highway  
567 BR-319 as a spearhead for Amazonian deforestation. *Land Use Policy*, 108, art. 105559.  
568 DOI: <https://doi.org/10.1016/j.landusepol.2020.104548>



- 569 Ferrante L, Fearnside PM (2018) Amazon sugarcane: A threat to the forest. *Science*,  
570 359, 1472. DOI: <https://doi.org/10.1126.aat4208>  
571
- 572 Ferrante L, Fearnside PM (2019) Brazil's new president and 'ruralists' threaten  
573 Amazonia's environment, traditional peoples and the global climate. *Environmental*  
574 *Conservation*, 46, 261-263. DOI: <https://doi.org/10.1017/S0376892919000213> .  
575
- 576 Ferrante L, Fearnside PM (2020a) Evidence of mutagenic and lethal effects of  
577 herbicides on Amazonian frogs. *Act Amazonica*, 50, 363-366. DOI:  
578 <https://doi.org/10.1590/1809-4392202000562>  
579
- 580 Ferrante L, Fearnside PM (2020b). The Amazon: biofuels plan will drive deforestation.  
581 *Nature*, 577, 170. DOI: <https://doi.org/10.1038/d41586-020-00005-8>  
582
- 583 Ferrante L, Fearnside PM (2021) Brazil's political upset threatens Amazonia. *Science*,  
584 371, 898. DOI: <https://doi.org/10.1126/science.abg9786>  
585
- 586 Ferrante L, Fearnside PM (2022a). Mining and Brazil's Indigenous peoples. *Science*,  
587 375, 276. DOI: <https://doi.org/10.1126/science.abn6753>  
588
- 589 Ferrante L, Fearnside PM (2022b) Countries should boycott Brazil over export-driven  
590 deforestation. *Nature*, 601, 318. DOI: <https://doi.org/10.1038/d41586-022-00094-7>  
591
- 592 Ferrante L, Fearnside PM (2022c) Indigenous lands protect Brazil's agribusiness.  
593 *Science*, 376, 810. DOI: <https://doi.org/10.1126/science.abq7243>  
594
- 595 Ferrante L, Fearnside PM (2023) Amazonia and the end of fossil fuels. *Nature*, 614,  
596 624. DOI: <https://doi.org/10.1038/d41586-023-00483-6>  
597
- 598 Ferrante L, Getirana A, Baccaro FB, Schöngart J, Leonel ACM, Gaiga R, Garey MV,  
599 Fearnside PM (2023) Effects of Amazonian flying rivers on frog biodiversity and  
600 populations in the Atlantic Rainforest. *Conservation Biology*, 37(3), art. e14033. DOI:  
601 <https://doi.org/10.1111/cobi.14033>  
602
- 603 Ferrante L, Leonel ACM, Gaiga R, Kaefer IL, Fearnside PM (2019) Local extinction of  
604 *Scinax caldarum*, a treefrog in Brazil's Atlantic Forest. *Herpetological Journal*, 29, 295-  
605 298. DOI: <https://doi.org/10.33256/hj29.4.295298>  
606
- 607 Ferrante L, Kaefer IL, Baccaro FB (2020) Aliens in the backyard: Did the American  
608 bullfrog conquer the habitat of native frogs in the semi-deciduous Atlantic Forest?  
609 *Herpetological Journal*, 30, 93-98. DOI: <https://doi.org/10.33256/hj30.2.9398>  
610
- 611 Ferreira M, Aleixo A, Ribas CC, Santos MPD (2016) Biogeography of the Neotropical  
612 genus *Malacoptila* (Aves: Bucconidae): The influence of the Andean orogeny,  
613 Amazonian drainage evolution and palaeoclimate. *Journal of Biogeography*, 44, 748-  
614 759. <https://doi.org/10.1111/jbi.12888>  
615
- 616 Flores BM, Holmgren M (2021) White-sand savannas expand at the core of the Amazon  
617 after forest wildfires. *Ecosystems*, 24, 1624–1637. DOI: <https://doi.org/10.1007/s10021-021-00607-x>  
618

- 619  
620 Gatti LV, Cunha CL, Marani L. et al. (2023) Increased Amazon carbon emissions  
621 mainly from decline in law enforcement. *Nature*, 621, 318–323. DOI:  
622 <https://doi.org/10.1038/s41586-023-06390-0>  
623
- 624 Gotelli NJ, Ellison AM (2004) A primer of ecological statistics. Sunderland, MA, USA,  
625 515 pp.  
626
- 627 Heyer WR, Rand AS, Cruz CAG, Peixoto OL (1988) Decimations, extinctions, and  
628 colonizations of frog populations in southeast Brazil and their evolutionary  
629 implications. *Biotropica*, 20, 230-235. DOI: <https://doi.org/10.2307/2388238>  
630
- 631 Hu J, Huang Y, Jiang J, Guisan A (2019) Genetic diversity in frogs linked to past and  
632 future climate changes on the roof of the world. *Journal of Animal Ecology*, 88, 953-  
633 963. <https://doi.org/10.1111/1365-2656.12974>  
634
- 635 INMET (Instituto Nacional de Meteorologia) (2023). Dados Históricos Anuais.  
636 <https://portal.inmet.gov.br/dadoshistoricos>  
637
- 638 IPCC (Intergovernmental Panel on Climate Change) (2014) Climate Change 2014:  
639 Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of  
640 Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on  
641 Climate Change. Barros VR et al. (eds.). Cambridge University Press, Cambridge, UK  
642 and New York, NY, USA. 688 pp. <https://www.ipcc.ch/report/ar5/wg2/>  
643
- 644 IPCC (Intergovernmental Panel on Climate Change) (2022) Climate Change 2022:  
645 Impacts, Adaptation and Vulnerability. Cambridge University Press, Cambridge, UK  
646 and New York, NY, USA Available at: [https://reliefweb.int/report/world/climate-  
647 change-2022-impacts-adaptation-and-  
648 vulnerability?gclid=Cj0KCQjwx5qoBhDyARIsAPbMagB68QNOJKKGRXrHIE-  
649 4LsOHeH-Ca70vPDKfvXfXzjk7MrDnEUfE86caAr9qEALw\\_wcB](https://reliefweb.int/report/world/climate-change-2022-impacts-adaptation-and-vulnerability?gclid=Cj0KCQjwx5qoBhDyARIsAPbMagB68QNOJKKGRXrHIE-4LsOHeH-Ca70vPDKfvXfXzjk7MrDnEUfE86caAr9qEALw_wcB)  
650
- 651 Joly CA, Padgurschi MCG, Pires APF et al. (2019) Apresentando o Diagnóstico  
652 Brasileiro de Biodiversidade e Serviços Ecossistêmicos. In: Joly CA, Scarano FR,  
653 Seixas CS et al. (eds.) 1º Diagnóstico Brasileiro de Biodiversidade e Serviços  
654 Ecossistêmicos. Editora Cubo, São Carlos, SP, Brazil. pp. 6-33.  
655 <https://www.bpbes.net.br/produto/diagnostico-brasileiro/>  
656
- 657 Jorge RF, Ferrão M, Lima AP (2020) Out of bound: A new threatened harlequin toad  
658 (Bufonidae, *Atelopus*) from the outer borders of the Guiana Shield in central Amazonia  
659 described through integrative taxonomy. *Diversity*, 12, art. 310. DOI:  
660 <https://doi.org/10.3390/d12080310>  
661
- 662 Kay G, Dunstone NJ, Smith DM, Betts RA, Cunningham C, Scaife AA (2022)  
663 Assessing the chance of unprecedented dry conditions over North Brazil during El Niño  
664 events. *Environmental Research Letters*, 17, art. 064016. DOI:  
665 <https://doi.org/10.1088/1748-9326/ac6df9>  
666
- 667 Lima AP, Magnusson WE, Menin M, Erdtmann LK, Rodrigues DJ, Keller C, Holdi W  
668 (2006) Guia de sapos da Reserva Adolpho Ducke, Amazônia Central = Guide to the

- 669 frogs of Reserva Adolpho Ducke, Central Amazonia. Áttema Design Editorial, Manaus,  
670 AM, Brazil. 168 pp.
- 671
- 672 Longhini LS, Zena LA, Polymeropoulos ET, Rocha ACG, da Silva LG, Prado CPA,  
673 Bícego KC, Gargaglioni LH (2021) Thermal acclimation to the highest natural ambient  
674 temperature compromises physiological performance in tadpoles of a stream breeding  
675 savanna tree frog. *Frontiers in Physiology*, 12, art. 726440. DOI:  
676 <https://doi.org/10.3389/fphys.2021.726440>
- 677
- 678 Loyola RD, Lemes P, Brum FT, Provete DB, Duarte LDS (2013) Clade-specific  
679 consequences of climate change to amphibians in Atlantic Forest protected areas.  
680 *Ecography*, 37, 65-72. DOI: <https://doi.org/10.1111/j.1600-0587.2013.00396.x>
- 681
- 682 Luenberger DG (1997) [1969] "Least-Squares Estimation". Optimization by Vector  
683 Space Methods. John Wiley & Sons, New York, NY, USA. pp. 78–102.
- 684
- 685 Magnusson WE, Mourão G (2005) Statistics without Math. Editora Planta / Sinauer  
686 Associates, Londrina, PR, Brazil. 136 pp.
- 687
- 688 Magnusson WE, Lima AP, Luizão RC, Luizão F, Costa FRC, Castilho CV, Kinupp VF  
689 (2005) RAPELD: Uma modificação do método de Gentry para inventários de  
690 biodiversidade em sítios para pesquisa ecológica de longa duração. *Biota Neotropica*,  
691 5(2), art: bn01005022005. DOI: <https://doi.org/10.1590/S1676-06032005000300002>
- 692
- 693 Magnusson WE, Pezzini RB, Baccaro FB, et al. (2013) Biodiversidade e  
694 Monitoramento Ambiental Integrado: O Sistema RAPELD na Amazônia. Attema  
695 Editorial, Santo André, SP, Brazil. 335 pp.
- 696
- 697 Magnusson WE (2020) Análise do EIA-RIMA do trecho do meio da rodovia BR-319.  
698 Re: Ofício nº 205/2019/9º OFÍCIO/PR/AM (expediente PR/AM-00055261/2019).  
699 <https://bit.ly/3LsA6zV>
- 700
- 701 Magrin GO, Marengo JA, Boulanger J-P, Buckeridge MS, Castellanos E, Poveda G,  
702 Scarano FR, Vicuña S (2014) Central and South America. In: Climate Change 2014:  
703 Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of  
704 Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on  
705 Climate Change. Barros VR et al. (eds). Cambridge University Press, Cambridge, UK  
706 and New York, NY, USA, pp. 1499-1566. <https://bit.ly/3TkeBFp>
- 707
- 708 Magurran AE (2004) Measuring Biological Diversity. Wiley-Blackwell, Hoboken, NJ,  
709 USA. 261 pp.
- 710
- 711 Marengo JA, Espinoza JC, Fu R, Muñoz JCJ, Alves LM, Rocha HR, Schöngart J (2021)  
712 Chapter 22: Long-term variability, extremes and changes in temperature and hydro  
713 meteorology in the Amazon region. In: Nobre C, Encalada A et al. (eds). Amazon  
714 Assessment Report 2021. United Nations Sustainable Development Solutions Network,  
715 New York, NY, USA. DOI: <https://doi.org/10.55161/ZGJG8060>
- 716
- 717 Menin M, Waldez F, Lima A (2008) Temporal variation in the abundance and number  
718 of species of frogs in 10,000 ha of a forest in central Amazonia, Brazil. *South*

- 719 *American Journal of Herpetology*, 3(1), 68-81. DOI: [https://doi.org/10.2994/1808-](https://doi.org/10.2994/1808-9798(2008)3[68:TVITAA]2.0.CO;2)  
720 9798(2008)3[68:TVITAA]2.0.CO;2  
721
- 722 Mills LS (2013) Conservation of Wildlife Populations: Demography, Genetics, and  
723 Management. John Wiley & Sons., Chichester, UK. 344 pp.  
724
- 725 Moraes LJCL, Pavan D, Barros MC, Ribas CC (2016) The combined influence of  
726 riverine barriers and flooding gradients on biogeographical patterns for amphibians and  
727 squamates in south-eastern Amazonia. *Journal of Biogeography*, 43, 2085–2309. DOI:  
728 <https://doi.org/10.1111/jbi.12756>  
729
- 730 Najjar T, Ferrante L (2018) The journey of life of the tiger-striped leaf frog *Callimedusa*  
731 *tomopterna* (Cope, 1868): Notes of sexual behaviour, nesting and reproduction in the  
732 Brazilian Amazon. *Herpetology Notes*, 11, 531-538.  
733 <https://repositorio.inpa.gov.br/bitstream/1/15326/1/artigo-inpa.pdf>  
734
- 735 Oliveira MD, Baccaro FB, Braga-Neto R, Magnusson WE (2008) Reserva Ducke: A  
736 Biodiversidade Amazônica através de uma Grade. PPBio. Áttema Design Editorial,  
737 Manaus, AM, Brazil. 166 pp. <https://bit.ly/3yC8vIO>  
738
- 739 Olson DM, Dinerstein E, Wikramanayake ED, Burgess ND, Powell GVN, Underwood  
740 EC, D'amico JA, Itoua I, Strand HE, Morrison JC, Loucks CJ, Allnutt TF, Ricketts TH,  
741 Kura Y, Lamoreux JF, Wettengel WW, Hedao P, Kassem KR (2001) Terrestrial  
742 Ecoregions of the World: A New Map of Life on Earth: A new global map of terrestrial  
743 ecoregions provides an innovative tool for conserving biodiversity. *BioScience*, 51:  
744 933–938. DOI: [https://doi.org/10.1641/0006-3568\(2001\)051\[0933:TEOTWA\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2001)051[0933:TEOTWA]2.0.CO;2)  
745
- 746 Paredes-Trejo F, Barbosa HA, Giovannettone J, Kumar TVL, Thakur MK, Buriti CA  
747 (2021). Long-term spatiotemporal variation of droughts in the Amazon River Basin.  
748 *Water* 13: art. 351. DOI: <https://doi.org/10.3390/w13030351>  
749
- 750 Pounds JA, Crump MI (1994) Amphibian declines and climate disturbance: the case of  
751 the golden toad and the harlequin frog. *Conservation Biology*, 8, 72–85 DOI:  
752 <http://www.jstor.org/stable/2386722>  
753
- 754 Pounds J, Fogden M, Campbell J (1999) Biological response to climate change on a  
755 tropical mountain. *Nature*, 398, 611–615. DOI: <https://doi.org/10.1038/19297>  
756
- 757 Quental TB, Marshall CR (2013) How the red queen drives terrestrial mammals to  
758 extinction. *Science*, 341, 290-292. DOI: <https://doi.org/10.1126/science.1239431>  
759
- 760 Ramos CJP, Graça PMLA, Fearnside PM (2018) Deforestation dynamics on an  
761 Amazonian peri-urban frontier: Simulating the influence of the Rio Negro Bridge in  
762 Manaus, Brazil. *Environmental Management*, 62(6), 1134-1149. DOI:  
763 <https://doi.org/10.1007/s00267-018-1097-3>  
764
- 765 Reis M, Graça PMLA, Yanai AM, Ramos CJP, Fearnside PM (2021) Forest fires and  
766 deforestation in the central Amazon: Effects of landscape and climate on spatial and  
767 temporal dynamics. *Journal of Environmental Management*, 288, art. 112310. DOI:  
768 <https://doi.org/10.1016/j.jenvman.2021.112310>

- 769  
770 Ribas CC, Aleixo A, Nogueira AC, Miyaki CY, Cracraft J (2012) A  
771 palaeobiogeographic model for biotic diversification within Amazonia over the past  
772 three million years. *Proceedings of the Royal Society B*, 279, 681–689. DOI:  
773 <https://doi.org/10.1098/rspb.2011.1120>  
774
- 775 Ribeiro MNG (1976) Aspectos climatológicos de Manaus. *Acta Amazonica*, 6(2), 229-  
776 233. DOI: <https://doi.org/10.1590/1809-43921976062229>  
777
- 778 Ritchie J, Dowlatabadi, H (2017) Why do climate change scenarios return to coal?  
779 *Energy*, 140, 1276-1291. DOI: <https://doi.org/10.1016/j.energy.2017.08.083>  
780
- 781 Rojas-Ahumada DP, Menin M (2010) Composition and abundance of Anurans in  
782 Riparian and Non-riparian Areas in A Forest in Central Amazonia, Brazil. *South*  
783 *American Journal of Herpetology*, 5(2), 157-167. DOI:  
784 <https://doi.org/10.2994/057.005.0210>  
785
- 786 Ruaro R, Ferrante L, Fearnside PM (2021) Brazil's doomed environmental licensing.  
787 *Science*, 372, 1049-1050. DOI: <https://doi.org/10.1126/science.abj4924>  
788
- 789 Ruaro R, Alves G, Tonella L, Ferrante L, Fearnside PM (2022) Loosening of  
790 environmental licensing threatens Brazilian biodiversity and sustainability. *Die Erde*,  
791 153, 60-64. DOI: <https://doi.org/10.12854/erde-2022-614>  
792
- 793 Sales LP, Galetti M, Pires MM (2020) Climate and land-use change will lead to a faunal  
794 “savannization” on tropical rainforests. *Global Change Biology*, 26, 7036-7044. DOI:  
795 <https://doi.org/10.1111/gcb.15374>  
796
- 797 Santos YLF, Yanai AM, Ramos CJP, Graça PMLA, Veiga JAP, Correia FWS,  
798 Fearnside PM (2022) Amazon deforestation and urban expansion: Simulating future  
799 growth in the Manaus Metropolitan Region, Brazil. *Journal of Environmental*  
800 *Management*, 304, art. 114279. DOI: <https://doi.org/10.1016/j.jenvman.2021.114279>  
801
- 802 Science on a Sphere (2023) Dataset Catalog. <https://sos.noaa.gov/catalog/datasets/>  
803
- 804 Shapiro SS, Wilk MB (1965) An analysis of variance test for normality (complete  
805 samples). *Biometrika*, 52, 591–611.  
806
- 807 Statsoft (2007) Statistica (data analysis software system) version 8.0. Statsoft, Inc.,  
808 Tulsa, OK, USA.  
809
- 810 Stebbins RC, Cohen NW (1995) A Natural History of Amphibians. Princeton  
811 University Press, UK, 336 pp.  
812
- 813 Strona G, Bradshaw CJA (2022) Coextinctions dominate future vertebrate losses from  
814 climate and land use change. *Science Advances*, 8, art. eabn434. DOI:  
815 <https://doi.org/10.1126/sciadv.abn4345>  
816
- 817 Stuart SN, Hoffmann M, Chanson JS, Cox NA, Berridge RJ, Ramani P, Young BE  
818 (2008) Threatened Amphibians of the World. Ed. Lynx Edicions, IUCN, and

- 819 Conservation International, Barcelona, Spain, Gland, Switzerland and Arlington, VA,  
820 USA. 758 pp.  
821
- 822 Tang S, Li Y, Wang Y (2001) Simultaneous equations, error-in-variable models, and  
823 model integration in systems ecology. *Ecological Modelling*, 142(3), 285–294. DOI:  
824 [https://doi.org/10.1016/s0304-3800\(01\)00326-x](https://doi.org/10.1016/s0304-3800(01)00326-x)  
825
- 826 Urbina-Cardona JN, Olivares-Perez M, Reynoso VH (2006) Herpetofauna diversity and  
827 microenvironment correlates across the pasture-edge-interior gradient in tropical  
828 rainforest fragments in the region of Los Tuxtlas, Veracruz. *Biological Conservation*,  
829 132, 61–75. DOI: <https://doi.org/10.1016/j.biocon.2006.03.014>  
830
- 831 Vasconcelos TS, do Nascimento BTM, Prado VHM (2018) Expected impacts of climate  
832 change threaten the anuran diversity in the Brazilian hotspots. *Ecology and Evolution*,  
833 17(3), 7894–7906. DOI: <https://doi.org/10.1002/ece3.6313>  
834
- 835 Verdade VK, Valdujo PH, Carnaval AC, Schiesari L, Toledo LF, Mott T, Andrade GV,  
836 Eterovick PC, Menin M, Pimenta BVS, Nogueira C, Lisboa CS, de Paula CD, Silvano  
837 DL (2012) A leap further: The Brazilian Amphibian Conservation Action Plan. *Alytes*  
838 29: 28-43. *Alytes: International Journal of Batrachology*, 29(1-4), 27-42.  
839 <https://bit.ly/3SZ0Wnt>  
840
- 841 Vieira KS, Montenegro PFG, Santana GG, Vieira WLDS (2018) Effect of climate  
842 change on distribution of species of common horned frogs in South America. *PLoS*  
843 *ONE*, 13(9), art. e0202813. DOI: <https://doi.org/10.1371/journal.pone.0202813>  
844
- 845 Vilani RM, Ferrante L, Fearnside PM (2023) The first acts of Brazil’s new president:  
846 Lula’s new Amazon institutionalality. *Environmental Conservation*, 50(3), 148 – 151.  
847 DOI: <https://doi.org/10.1017/S0376892923000139>  
848
- 849 Weygoldt P (1989) Changes in the composition of mountain stream frog communities in  
850 the Atlantic mountains of Brazil frogs as indicators of environmental deteriorations.  
851 *Studies on Neotropical Fauna and Environment*, 24, 249-256. DOI:  
852 <https://doi.org/10.1080/01650528909360795>  
853
- 854 Zipkin EF, DiRenzo GV, Ray JM, Rossman S, Lips KR (2020) Tropical snake diversity  
855 collapses after widespread amphibian loss. *Science*, 367, 814–816. DOI:  
856 <https://doi.org/10.1126/science.aay5733>

# Supplementary Material

## Climate change in the Central Amazon and its impacts on frog populations

Lucas Ferrante<sup>1, 2,\*</sup>, Diana Rojas-Ahumada<sup>2</sup>, Marcelo Menin<sup>1,†</sup>, & Philip Martin Fearnside<sup>3</sup>

<sup>1</sup>Universidade Federal do Amazonas (UFAM), Manaus, Amazonas, Brazil.

<sup>2</sup>Programa de Pós-Graduação em Ecologia, Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Amazonas, Brazil.


<sup>3</sup>Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Amazonas, Brazil.

\*Correspondence to: lucasferrante@hotmail.com

† Posthumous author.

Appendix S1 Climate data.....	2
Appendix S2. Normality tests.....	3
Appendix S3. Homoscedasticity test.....	6

## Appendix S1 Climate data





## Appendix S2. Normality tests (Shapiro-Wilk test):

Shapiro-Wilk results:

*A. spumarinus*

W: 0.7478694

p-value: 0.02843795

*A. spumarinus* has explosive reproduction, which tends to interfere with the density of individuals and which explains the absence of normality for this species.

-----

*L. pentadactylus*

W: 0.8332792

p-value: 0.1471844

There is no evidence to reject the null hypothesis of normality.

-----

*O. oophagus*

W: 0.981019

p-value: 0.9399969

There is no evidence to reject the null hypothesis of normality.

-----

*L. rhondonmystax*

W: 0.8327445

p-value: 0.1458437

There is no evidence to reject the null hypothesis of normality.

-----

*P. fenestratus*

W: 0.925325

p-value: 0.5648562

There is no evidence to reject the null hypothesis of normality.

-----

*S. mirandariberoi*

W: 0.8767309

p-value: 0.2947585

There is no evidence to reject the null hypothesis of normality.

-----


*S. salseri*







W: 0.874595

p-value: 0.2855315

There is no evidence to reject the null hypothesis of normality.

# Shapiro-Wilk graphs:





### Appendix S3. Homoscedasticity test:

Largest set of species with homogeneity of variances: *L. pentadactylus* (n=63), *L. rhodomystax* (n=15), *O. oophagus* (n=3222), *P. fenestratus* (n=6702), *S. mirandariberoi* (n=1459) and *S. salseri* (n=996).

$p > 0.05$  (There is no evidence to reject homoscedasticity)

*L. pentadactylus*: Average: 12.60, Variance: 8.24.

*L. rhodomystax*: Average: 3.00, Variance: 1.20.

*O. oophagus*: Average: 644.40, Variance: 58124.24.

*P. fenestratus*: Average: 1340.40, Variance: 371489.04.

*S. mirandariberoi*: Average: 291.80, Variance: 58563.76.

*S. salseri*: Average: 199.20, Variance: 26727.76.

\* *A. manauensis* does not show homoscedasticity in relation to the other species; these results may have been influenced by the explosive reproduction habit of this species.

