

This file has been cleaned of potential threats.

If you confirm that the file is coming from a trusted source, you can send the following SHA-256 hash value to your admin for the original file.

0d1c846e34987d64e183465ebe74aa3205d81d228e9c982f70579a2a80b301c5

To view the reconstructed contents, please SCROLL DOWN to next page.

**The text that follows is a PREPRINT.  
O texto que segue é um PREPRINT.**

*Please cite as:*

*Favor citar como:*

Azevedo-Santos, V.M., P.M. Fearnside,  
M.S. Arcifa, L.H. Tonella, T.  
Giarrizzo, F.M. Pelicice, A.A.  
Agostinho, A.E. Magurran & N.L.  
Poff. 2024. **Irrigation dams  
threaten Brazilian  
biodiversity.** *Environmental  
Management* <https://doi.org/10.1007/s00267-024-01946-w> (in press).

ISSN: 0364-152X

DOI: 10.1007/s00267-024-01946-w

Copyright: Springer.

The original publication will be available at:  
A publicação original estará disponível em:

<https://doi.org/10.1007/s00267-024-01946-w>  
<https://link.springer.com/journal/267>

# Irrigation dams threaten Brazilian biodiversity

Valter M. Azevedo-Santos<sup>1,2\*</sup>, Philip M. Fearnside<sup>3</sup>, Marlene S. Arcifa<sup>4</sup>, Livia H. Tonella<sup>5</sup>, Tommaso Giarrizzo<sup>2,6</sup>, Fernando M. Pelicice<sup>1,7</sup>, Angelo A. Agostinho<sup>8</sup>, Anne E. Magurran<sup>9</sup>, N. LeRoy Poff<sup>10</sup>

<sup>1</sup>Universidade Federal do Tocantins (UFT), Programa de Pós-Graduação em Biodiversidade, Ecologia e Conservação, Porto Nacional CEP 77500-000, Tocantins, Brazil.

<sup>2</sup>Grupo de Ecologia Aquática, Espaço Inovação do Parque de Ciência e Tecnologia Guamá (PCT Guamá), Belém, Pará, Brazil

<sup>3</sup>Instituto Nacional de Pesquisas da Amazônia (INPA), Av. André Araújo, 2936, 69067-375, Manaus, Amazonas, Brazil.

<sup>4</sup>Departamento de Biologia, Universidade de São Paulo, Av. Bandeirantes, 3900, Ribeirão Preto 14040-901, São Paulo, Brazil.

<sup>5</sup>Departamento de Direito, Universidade Federal do Tocantins (UFT), Palmas, Tocantins, Brazil.

<sup>6</sup>Instituto de Ciências do Mar (LABOMAR), Universidade Federal do Ceará (UFC), Avenida da Abolição 3207, Fortaleza 60165-081, Ceará, Brazil.

<sup>7</sup>Universidade Federal do Tocantins, Núcleo de Estudos Ambientais, 77500-000 Porto Nacional, TO, Brazil.

<sup>8</sup>Programa de Pós-Graduação em Ecologia de Ambientes Aquáticos Continentais (PEA), Universidade Estadual de Maringá, PR, Brazil.

<sup>9</sup>Centre for Biological Diversity, School of Biology, University of St Andrews, Scotland.

<sup>10</sup>Graduate Degree Program in Ecology, Colorado State University, Fort Collins CO 80524 USA, and Centre for Applied Water Science, University of Canberra, ACT 2617 Australia.

\*Corresponding author (E-mail: valter.ecologia@gmail.com)

## Acknowledgements

PMF was supported by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) (2020/08916-8), Fundação de Amparo à Pesquisa do Estado do Amazonas (FAPEAM) (0102016301000289/2021-33), FINEP/Rede CLIMA (01.13.0353-00). The Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) provided research grants to PMF (Process 312450/2021-4), FMP (Process 312256/2020-5), and AAA (Process 312549/2021). Leverhulme Trust (RPG- 2019-402) provided research grants to AEM.

## Declarations

### Funding

PMF was supported by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) (2020/08916-8), Fundação de Amparo à Pesquisa do Estado do Amazonas (FAPEAM) (0102016301000289/2021-33), FINEP/Rede CLIMA (01.13.0353-00). The Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) provided research grants to PMF (Process 312450/2021-4), FMP (Process 312256/2020-5), and

AAA (Process 312549/2021). Leverhulme Trust (RPG- 2019-402) provided research grants to AEM.

Conflicts of interest/Competing interest  
None.

Availability of data and material  
Not applicable.

Code availability  
Not applicable

Authors' contributions  
Valter M. Azevedo-Santos conceived the idea of the article and wrote the first draft. All authors contributed with ideas, text, and corrections. All authors approved the present version of the article.

Ethics approval (include appropriate approvals or waivers)  
Not applicable.

Consent to participate  
All authors consent with the content of the paper.

Consent for publication (include appropriate statements)  
All authors consent with the publication of the paper

### **Abstract**

Brazil is among the main contributors to global biodiversity, which, in turn, provides enormous ecosystem services. Agriculture is one of the activities most benefited by these ecosystem services. However, this activity has contributed to the degradation of aquatic and terrestrial ecosystems and eroding Brazilian biodiversity. This conflict is growing, as emerging unsustainable legislative proposals to benefit the agricultural sector are likely to increase the decline of biodiversity. For instance, one such initiative (Bill 1282/2019) intends to change the “Forest Code” (Law 12,651/2012) to facilitate the construction of irrigation dams in Permanent Protected Areas; this last includes strips (with vegetation or no) along the edges of watercourses. In addition, two other similar bills are under progress in the Chamber of Deputies. Here we detailed these three bills and discuss — if approved — their consequences for Brazil’s biodiversity. Expected negative impacts with changes in the legislation include, for example, increased deforestation, siltation, habitat fragmentation, introduction of non-native species, reductions in the availability of aquatic habitats, and changes in biogeochemical process. These proposals jeopardize biodiversity and may compromise the negotiations for an agreement between Mercosur and the European Union.

### **Keywords**

Agriculture; Deforestation; Forest Code; Global warming; Habitat fragmentation; Law

### **Introduction**

Brazil’s is among the main contributors to global biodiversity (Agostinho et al. 2005). Past estimates suggested that there are around 210,000 known species in the

country (Lewinsohn and Prado 2005) — but this number is currently believed to be greater. Brazil’s biodiversity provides numerous ecosystem services, including those classified as regulating, support, provision, and cultural (e.g., Ellwanger et al. 2022; Pelicice et al. 2023). This biodiversity has been increasingly threatened by the expansion of human activities (e.g., Loisel et al. 2010; Fernandes et al. 2016, 2020; Garcia et al. 2021), including through official policies (e.g., Fearnside 2016; Pelicice et al. 2017).

Agriculture activities have generated negative impacts on Brazilian environments and biodiversity (Hepp et al. 2010; Pelicice et al. 2021). In the last 50 years, the irrigated area in Brazil increased by approximately a factor of 10, and in 2015, this area totaled ~7 million hectares, with projections indicating it would reach 10 million hectares by 2030 (ANA 2019). Irrigation occurs in different systems (e.g., center-pivots, drip) with water taken from hydroelectric reservoirs, lakes and, also frequently, rivers and streams. Streams can also be dammed to supply water for irrigation, even though these dams are in several cases illegal and negatively affect biodiversity (Maffra and Souza 2018). The expansion of irrigated fields may accelerate even further with recent legislation favoring — in an unsustainable way — the agricultural sector.

Brazil’s current “Forest Code” (Law number 12,651/2012) requires the maintenance of Permanent Preservation Areas (known as “APPs” in Portuguese), which include forest strips along the edges of watercourses (Brazil 2012). Interventions in APPs are only allowed in specific cases when impacts are “low” (Brazil 2012). However, three bills currently advancing through committees in the Chambers of Deputies and the Senate would facilitate the construction of dams for irrigation in these areas. Here we explain why these proposed laws threaten biodiversity.

### **Facilitating the construction of irrigation dams**

The importance of riparian habitats is recognized by Brazil’s current “Forest Code” (Law 12,651/2012), which establishes requirements for the protection of vegetation along waterbodies, both lotic and lentic (Brazil 2012: Chapter II, Article 4°). For instance, the protection of a 30-meter-wide strip is required along each side of a stream with a width of 9 meters (Table 1). Law 12,651/2012 clearly states that deforestation in the Permanent Preservation Areas (hereafter APPs) may be done only in cases of need for “(...) public utility, social interest or with low environmental impact as foreseen in this Law” (Brazil 2012: Chapter II, Article 8). This limitation may change with the pending bills.

Three bills (two in the Chamber of Deputies and one in the Senate) have been proposed to change Law 12,651/2012 to allow interventions in APPs. Bill 2168/2021 (Chamber of Deputies) proposes a change “(...) to consider irrigation and animal watering infrastructure as public utility works, including dams or the damming of watercourses that cause intervention or suppression of native vegetation in a permanent preservation area (...)” (Brazil 2021a: 1). In turn, Bill 399/2022 (Chamber of Deputies) aims “(...) to turn the areas destined for irrigation into public utilities and social interest” (Brazil 2022: 1), thus allowing irrigation dams to be built without regard for impacts and degradation of APPs. Lastly, Bill 1282/2019 (Senate) proposes the following provisions: “In rural properties is allowed (...) the construction of water reservoirs for irrigation projects and the physical infrastructure associated with them” (Brazil 2019a: 2). The justification of the bill states that “(...) the lack of clarity in the current legislations on the subject has been, for a long time, hindering the expansion of technologies related to irrigation” (Brazil 2019a: 3), and that “(...) the insertion of this

device in the new Brazilian Forest Code will bring clarity necessary for a topic of such significance and fundamental importance for the reduction of crop losses (...)” (Brazil 2019a: 3). The reference to “water reservoirs” means that they include the riparian region (adjacent to the waterbody) that would be flooded.

**Table 1.** Size of Permanent Preservation Areas according to the Article 4° of Law 12,651/2012 (only Items I and II) (Brazil 2012).

Type of waterbody	Item in Law 12,651	Width or size of the waterbody	Width of the lateral strip protected (meters)
Watercourse	I	<10 wide	30
Watercourse	I	Between 10 to 50 wide	50
Watercourse	I	Between 50 to 200 wide	100
Watercourse	I	Between 200 to 600 wide	200
Watercourse	I	> 600 wide	500
Lake or lagoon	II	Any size	100 (rural areas)
Lake or lagoon	II	Up to 20 ha of surface	50 (rural areas)
Lake or lagoon	II	Any size	30 (urban areas)

Note: The width of the lateral strip protected along the watercourse (item I) may be lower in urban areas, depending on the local administration (see Azevedo-Santos et al. 2023)

These proposals can enable the degradation or the suppression of riparian forest areas by simplifying the process of dam building for irrigation — especially in streams and small rivers. This implies that interventions in APPs will occur without rigorous authorization procedures or technical assessments. These three bills, once approved, have the potential to cause a significant setback in the scenario of environmental policies and management of freshwater resources, watercourses, and associated biodiversity.

### Threats to biodiversity

Facilitating the construction of dams in APPs will increase different negative impacts on aquatic and terrestrial ecosystems and their biodiversity. Below we will exemplify the expected negative impacts if any of these three bills is approved.

The most obvious consequence will be increased deforestation of APPs. Existing vegetation adjacent to the waterbody will be degraded or lost, either through direct removal or inundation created by the dam — as generally occurs (see Maffra and Souza 2018). Deforestation, beyond results in the loss of tree species, eliminates habitat for terrestrial organisms, such as insects, reptiles, birds, and mammals that live and depend on the riparian vegetation (Develey and Pongiluppi 2010; Freitas et al. 2010; Galetti et al. 2010; Marques et al. 2010; Farias et al. 2015). In addition, the removal of riparian vegetation can trigger siltation processes in aquatic ecosystems (Mittermeier et al. 1990), a phenomenon with adverse consequences for macroinvertebrates and fish (Casatti 2004; Couceiro et al. 2010). In general, the removal of riparian vegetation can also harm fish populations and assemblages (Azevedo-Santos et al. 2021) and affect biotic interactions associated with seed dispersion (Nogueira et al. 2023).

Facilitating infrastructure to block watercourses will increase habitat fragmentation. Dams, including small irrigation dams, impede the free movement of fish and invertebrates (Poff and Hart 2002; Pelicice et al. 2015; Chappell et al. 2019), affecting also drift dispersion of animal and plant propagules (Brooks et al. 2018). This phenomenon, which is documented in watercourses with barriers (e.g., Agostinho et al.

2008; Brejão et al. 2020), can affect gene flow, dispersion (including migration), and demographic dynamics.

The introduction of non-native species will also be intensified, especially considering the Brazilian scenario. Dams built to supply water, including for irrigation, have been — and may be — used for fish production in some cases (Albinati 2006; Cardoso Filho et al. 2010). Small dams used for recreational fishing and aquaculture have been a regular source of non-native species in Brazil (Vitule et al. 2006; Forneck et al. 2021). Moreover, there is a growing incentive for raising non-native fish in Brazil (e.g., Coelho and Henry 2017; Brito et al. 2018; Latini et al. 2021). The proliferation of dams, therefore, must be accompanied by the spread of non-native organism. We highlight that introduction of non-native species — including freshwater fishes — is a threat to native species (Vitule et al. 2009; Simberloff et al. 2023).

Other expected increases in negative effects relate to water use. Removing water from watercourses (either for agriculture or for urban use) reduces the availability of habitat and increases the frequency of low-water conditions; and this cause shifts in the presence and abundance of species, including invertebrates and fishes (Walters and Post 2011; Grantham et al. 2013). Small dams cause similar low-water downstream effects that negatively affect aquatic species (Couto et al. 2023).

Lastly, changes in biogeochemical processes can be expected to increase if any of the three bills is approved. The alteration of flows due to dams can also modify the concentration of nutrients such as phosphorus and nitrogen, accumulating in the impoundment area and generating eutrophication events (Maavara et al. 2020). Such nutrient buildups reduce dissolved oxygen levels, affecting aquatic communities (Wurtsbaugh et al. 2019). In addition, deforestation on the border of created reservoirs reduce the quality of water (with the input solids and iron) used for irrigation (Favero et al. 2022), as well as compromise downstream habitats. Finally, reservoirs are known to be a source of greenhouse gas (Fearnside 2002; Maavara et al. 2020). These compounds lead to global warming (Fearnside 2002), consequently affecting biodiversity at different scales (Malcolm et al. 2006; Pounds et al. 2006).

The intensification of the above negative impacts can be expected to result in biodiversity losses, especially in a context where numerous Brazilian species already are threatened with extinction (e.g., MMA 2022). For example, a total of 311 fish species have been listed as threatened, and habitat loss and fragmentation, also associated with river damming, have been the main stressors (Santana et al. 2021). In addition, threatened fish species (e.g., '*Chasmocranus brachynema* Gomes & Schubart, 1958) (see Deprá and Slobodian 2024) do not have any of their habitat protected. The number of threatened species can be expected to grow with the proliferation of new dams, especially because many species are only known to occur in limited areas that are being converted to agriculture and ranching (e.g., Costa et al. 2022). In the case of fish, various species were only described in the last few years (e.g., Burger et al. 2019; Guimarães et al. 2019; Costa and Katz 2021; Deprá et al. 2022), indicating that the number of fish species present is even higher, especially for groups that also inhabit streams (e.g., characids, heptapterids, trichomycterids). The number of undescribed species of insects that live in riparian habitats is probably even more underestimated. New species are being described for plants in areas subjected to agriculture expansion (e.g., Carvalho et al. 2023), indicating that some may have been lost with no scientific record. The degradation of riparian habitats in the Brazilian Amazon, a megadiverse region, is especially concerning. This scenario stresses the need to avoid unsustainable policies like those discussed herein.

**Final remarks**

Various legislative proposals have emerged to weaken Brazil's current "Forest Code" (Law 12,651/2012), which would endanger Brazil's ecosystems (Table 2). These proposals include the three bills to remove restrictions on irrigation dams. All these bills must be defeated in the Brazilian National Congress and the discussions about them must include the participation of scientists (Azevedo-Santos et al. 2017), including experts in water resources management. We recognize that the recent increase in the representation of agribusiness interests ("ruralists") in Brazil's National Congress (Fearnside 2023) is a powerful barrier to this.



**Table 2.** Proposed law to make changes of the Permanent Preservation Areas of the “Forest Code” (Law 12,651/2012) and their expected main negative impacts mainly on Brazil’s ecosystems.

Bill	Legislative chamber	Situation	Description	Changes	Examples of expected negative impacts
Proposed Law 5650/2016	Chamber of Deputies	In progress	“Authorizes the planting of yerba mate in a Permanent Preservation Area on small properties or family rural possessions” (Brazil 2016).	Allows the farming of <i>Ilex paraguariensis</i> in Permanent Preservation Areas	In the cultivation of <i>Ilex paraguariensis</i> , producers may use fertilizing with nitrogen and phosphorus (e.g., Alves et al. 2017). In addition, there are human activities (e.g., harvest) in an area which will have majority a single species (i.e., yerba mate). Main expected negative effects are the pollution of waterbodies (with nutrients), stomping on non-target vegetation and habitat simplification.
Proposed Law 7592/2017	Chamber of Deputies	Suspended	“Amends Law No. 12,651, of May 25, 2012, to include the construction of hospitals and schools among the activities considered to be of social interest for the purpose of suppressing the Permanent Preservation Area” (Brazil 2017).	Allows intervention (removing vegetation is included) in Permanent Preservation Areas.	Intervention in riparian regions results in deforestation, erosion and siltation, pollution, increased brightness in watercourses (Mittermeier et al. 1990; Thomaz et al. 2021; Ottoni et al. 2023). These negative effects are expected to increase if the Bill is unarchived and approved.
Proposed Law 1.282/2019	Senate	In progress	“Amends the Forest Code to allow, in areas of permanent preservation of rural properties, the construction of water reservoirs for irrigation projects and the physical infrastructure associated with it” (Brazil 2019a).	Facilitate the construction of dams for irrigation within the APPs.	See section “Threats to biodiversity” of this article.
Proposed Law 1731/2019	Senate	In progress	“Amends Law No. 12,651, of 25 May 2012, which provides for the protection of native vegetation and other measures, to deal with	Allow agriculture and the presence of houses within	Intervention in riparian regions results in deforestation, erosion and siltation, pollution, increased brightness in watercourses (Mittermeier et al. 1990; Thomaz et al. 2021; Ottoni et al. 2023).

			rules on the occupation of Permanent Preservation Areas (APP)” (Brazil 2019b).	the limits of a Permanent	These negative effects are expected to increase if the Bill is approved.
Proposed Law 2510/2019	Chamber of Deputies	Approved - Now Law 14.285/2021	“Amends Law No. 12,651, of 25 May 2012, to provide for permanent protection areas in the urban perimeter and in metropolitan regions” (Brazil 2019c).	Allows to reduce the width of Permanent Preservation Area in the urban perimeter	Intervention in riparian regions results in deforestation, erosion and siltation, pollution, increased brightness in watercourses (Mittermeier et al. 1990; Thomaz et al. 2021; Ottoni et al. 2023). These negative effects are expected to increase with the approved law 14.285/2021 (Azevedo-Santos et al. 2023).
Proposed Law 2168/21	Chamber of deputies	In progress	“Amends Law 12,651/2012, to consider irrigation infrastructure works and animal watering as of public utility” (Brazil 2021a).	Facilitate the construction of dams for irrigation and for animal watering within Permanent Preservation Areas	See section “Threats to biodiversity” of this article. Areas used to livestock are subjected to fecal pollution (Souza et al. 2006). Therefore, this is also an expected negative impact if the Bill is approved.
Proposed Law 399/2022	Chamber of deputies	In progress	“Amends Law No. 12,651, of 25 May 2012, to make areas intended for irrigation public utility and social interest” (Brazil 2022).	Facilitate the construction of dams for irrigation	See section “Threats to biodiversity” of this article.

Note: The suspension of processing does not guarantee that processing will not be resumed or that another similar law will not be proposed.

Opposing irrigation dams in APPs does not mean being against agricultural expansion, but rather taking care not to exacerbate the biodiversity crisis. The three bills removing key restrictions on irrigation dams are not compatible with minimum levels of sustainability in agriculture. This is an issue of increasing concern among Brazil's international trading partners and is currently being raised in negotiations for an agreement between Mercosur and the European Union. Defeating these bills would be an important step towards addressing this challenge. It also underlines the need for Brazilian lawmakers to radically change their outlook on questions of sustainability to ensure that Brazil's rich biodiversity — that benefit the world — is preserved.

## References

- Agostinho AA, Thomaz SM, Gomes LC (2005) Conservation of the biodiversity of Brazil's inland waters. *Conserv Biol* 19:646–652. <https://doi.org/10.1111/j.1523-1739.2005.00701.x>
- Agostinho AA, Pelicice FM, Gomes LC (2008) Dams and the fish fauna of the Neotropical region: impacts and management related to diversity and fisheries. *Braz J Biol* 68: 1119–1132. <https://doi.org/10.1590/S1519-69842008000500019>
- Albinati RCB (2006) Aquicultura em pequenos açudes no semiárido. *Bahia Agrícola* 7(2): 66–72.
- Alves MV, Rovani M, Delazeri P et al (2017) Desempenho nutricional da erva-mate com diferentes formas de adubação. *Varia Scientia Agrárias* 5(2): 93–107.
- ANA (2019) Levantamento da agricultura irrigada por pivôs centrais no Brasil (1985-2017). Levantamento da agricultura irrigada por pivôs centrais no Brasil / Agência Nacional de Águas, Embrapa Milho e Sorgo. - 2. ed. Agência Nacional de Águas. Brasília, DF. 47 pp.
- Azevedo-Santos VM, Fearnside PM, Oliveira CS et al (2017) Removing the abyss between conservation science and policy decisions in Brazil. *Biodivers Conserv* 26: 1745–1752. <https://doi.org/10.1007/s10531-017-1316-x>
- Azevedo-Santos VM, Arcifa MS, Brito MF et al (2021) Negative impacts of mining on Neotropical freshwater fishes. *Neotrop Ichthyol* 19: e210001. <https://doi.org/10.1590/1982-0224-2021-0001>
- Azevedo-Santos VM, Daga VS, Tonella LH et al (2023) Brazil's urban ecosystems threatened by law. *Land Use Policy* 131: 106721. <https://doi.org/10.1016/j.landusepol.2023.106721>
- Brazil (2012) “Forest Code” - Law nº 12.651/2012 ([https://www.planalto.gov.br/ccivil\\_03/\\_ato2011-2014/2012/lei/112651.htm](https://www.planalto.gov.br/ccivil_03/_ato2011-2014/2012/lei/112651.htm)). Retrieved on 2 June 2023.
- Brazil (2016) Bill 5650/2016 (<https://www.camara.leg.br/proposicoesWeb/fichadetramitacao?idProposicao=2089024>). Retrieved on 2 June 2023.
- Brazil (2017) Bill 7592/2017 (<https://www.camara.leg.br/proposicoesWeb/fichadetramitacao?idProposicao=2136491>). Retrieved on 2 June 2023.
- Brazil (2019a) Bill 1282/2019 (<https://www25.senado.leg.br/web/atividade/materias/-/materia/135567>). Retrieved on 14 August 2023.
- Brazil (2019b) Bill 1731/2019 (<https://www25.senado.leg.br/web/atividade/materias/-/materia/135948>). Retrieved on 14 August 2023.

- Brazil (2019c) Bill 2510/2019 (<https://www.camara.leg.br/proposicoesWeb/fichadetramitacao?idProposicao=2199215>). Retrieved on 14 August 2023.
- Brazil (2021a) Bill 2168/2021 (<https://www.camara.leg.br/propostas-legislativas/2286627>). Retrieved on 14 August 2023.
- Brazil (2022) Bill 399/2022 (<https://www.camara.leg.br/propostas-legislativas/2316569>). Retrieved on 14 August 2023.
- Brejão GL, Teresa FB, Gerhard P (2020) When roads cross streams: fish assemblage responses to fluvial fragmentation in lowland Amazonian streams. *Neotrop Ichthyol* 18(3): e200031. <https://doi.org/10.1590/1982-0224-2020-003>
- Brito MFG, Magalhães ALB, Lima-Junior DP et al (2018). Brazil naturalizes non-native species. *Science* 361(6398): 139–139. <https://www.science.org/doi/10.1126/science.aau3368>
- Brooks AJ, Wolfenden B, Downes BJ, Lancaster J (2018). Barriers to dispersal: The effect of a weir on stream insect drift. *River Res Appl* 34(10): 1244–1253. <https://doi.org/10.1002/rra.3377>
- Burger R, Carvalho FR, Zanata AM (2019) A new species of *Astyanax* Baird & Girard (Characiformes: Characidae) from western Chapada Diamantina, Bahia, Brazil. *Zootaxa* 4604(2): 369–380. <https://doi.org/10.11646/zootaxa.4604.2.9>
- Cardoso Filho R, Campeche DFB, Paulino RV (2010) Tilápia em reservatório de água para irrigação e avaliação da qualidade da água. *Rev Bras Cienc Agrar* 5: 117–122. <http://www.redalyc.org/articulo.oa?id=119012589019>
- Carvalho CS, Lima HC, Zuanny DC et al (2023) The discovery of a new giant legume tree species in a severely fragmented landscape underscores the alarming threats to the biodiversity of the Brazilian Atlantic Forest. *Bot J Linn Soc* 201(2): 215–229. <https://doi.org/10.1093/botlinnean/boac034>
- Casatti L (2004) Ichthyofauna of two streams (silted and reference) in the upper Paraná river basin, southeastern Brazil. *Braz J Biol* 64: 757–765. <https://doi.org/10.1590/S1519-69842004000500004>
- Chappell J, McKay SK, Freeman MC, Pringle CM (2019) Long-term (37 years) impacts of low-head dams on freshwater shrimp habitat connectivity in northeastern Puerto Rico. *River Res Appl* 35(7): 1034–1043. <https://doi.org/10.1002/rra.3499>
- Coelho PN, Henry R (2017) The small foreigner: new laws will promote the introduction of non-native zooplankton in Brazilian aquatic environments. *Acta Limnol Bras* 29:e7. <https://doi.org/10.1590/S2179-975X0717>
- Costa WJEM, Katz AM (2021) Integrative taxonomy supports high species diversity of south-eastern Brazilian mountain catfishes of the *T. reinhardti* group (Siluriformes: Trichomycteridae). *Syst Biodivers* 19: 601–621. <https://doi.org/10.1080/14772000.2021.1900947>
- Costa WJEM, Feltrin CRM, Katz AM (2022) An endangered new catfish species of the genus *Cambeva* (*Cambeva gamabelardense* n. sp.) (Siluriformes, Trichomycteridae) from the Rio Chapecó drainage, southern Brazil. *Anim Biodivers Conserv* 45: 123–129. <https://doi.org/10.32800/abc.2022.45.0123>
- Couceiro SRM, Hamada N, Forsberg BR, Padovesi-Fonseca C (2010) Effects of anthropogenic silt on aquatic macroinvertebrates and abiotic variables in streams in the Brazilian Amazon. *J Soils Sediments* 10: 89–103. <https://doi.org/10.1007/s11368-009-0148-z>
- Couto TBA, Rezende RS, de Aquino PPU et al (2023) Effects of small hydropower dams on macroinvertebrate and fish assemblages in southern Brazil. *Freshw Biol* 68:956–971. <https://doi.org/10.1111/fwb.14078>

- Develey PF, Pongiluppi T (2010) Impactos potenciais na avifauna decorrentes das alterações propostas para o Código Florestal Brasileiro. *Biota Neotrop* 10: 43–45. <https://doi.org/10.1590/S1676-06032010000400005>
- Deprá GC, Aguilera G, Faustino-Fuster DR et al (2022) Redefinition of *Heptapterus* (Heptapteridae) and description of *Heptapterus carmelitanorum*, a new species from the upper Paraná River basin in Brazil. *Zoosyst Evol* 98: 327–343. <https://doi.org/10.3897/zse.98.89413>
- Deprá GC, Slobodian V (2024) Redescription of ‘*Chasmocranus*’ *brachynema* (Heptapteridae: Heptapterini). *Neotrop Ichthyol* (*in press*). <https://doi.org/10.1590/1982-0224-2023-0091>
- Ellwanger JH, Nobre CA, Chies JAB (2022) Brazilian biodiversity as a source of power and sustainable development: A neglected opportunity. *Sustainability* 15(1): 482. <https://doi.org/10.3390/su15010482>
- Farias TN, Rosa AHM, Damasceno PG et al (2015) Avifauna associada à Área de Preservação Permanente urbana de Tangará da Serra-MT. *Encicl Biosf* 11: 2945–2955.
- Favero D, Cotta AJB, Bonomo R, Rodrigues MB (2022) Hydrogeochemical characterization and suitability of water for irrigation in new and old reservoirs in northern Espírito Santo, Brazil. *Environ Dev Sustain* 24: 320–351. <https://doi.org/10.1007/s10668-021-01435-8>
- Fearnside PM (2002) Greenhouse gas emissions from a hydroelectric reservoir (Brazil’s Tucuruí Dam) and the energy policy implications. *Wat Air and Soil Poll* 133: 69–96. <https://doi.org/10.1023/A:1012971715668>
- Fearnside PM (2016) Brazilian politics threaten environmental policies. *Science* 353(6301): 746–748. <http://dx.doi.org/10.1126/science.aag0254>
- Fearnside PM (2023) The outlook for Brazil’s new presidential administration. *Trends Ecol Evol* 38(5): 387–388. <https://doi.org/10.1016/j.tree.2023.01.002>
- Fernandes GW, Goulart FF, Ranieri BD et al (2016) Deep into the mud: ecological and socio-economic impacts of the dam breach in Mariana, Brazil. *Nat Conserv* 14(2): 35–45. <http://dx.doi.org/10.1016/j.ncon.2016.10.003>
- Fernandes LFL, Paiva TRM, Longhini CM et al (2020) Marine zooplankton dynamics after a major mining dam rupture in the Doce River, southeastern Brazil: Rapid response to a changing environment. *Sci Total Environ* 736: 139621. <https://doi.org/10.1016/j.scitotenv.2020.139621>
- Forneck SC, Dutra FM, de Camargo MP et al (2021). Aquaculture facilities drive the introduction and establishment of non-native *Oreochromis niloticus* populations in Neotropical streams. *Hydrobiologia* 848: 1955–1966. <https://doi.org/10.1007/s10750-020-04430-8>
- Freitas AVL (2010) Impactos potenciais das mudanças propostas no Código Florestal Brasileiro sobre as borboletas. *Biota Neotrop* 10: 53–57. <https://doi.org/10.1590/S1676-06032010000400007>
- Galetti M, Pardini R, Duarte JMB et al (2010) Mudanças no Código Florestal e seu impacto na ecologia e diversidade dos mamíferos no Brasil. *Biota Neotrop* 10: 47–52. <https://doi.org/10.1590/S1676-06032010000400006>
- Garcia LC, Szabo JK, Roque FO et al (2021) Record-breaking wildfires in the world’s largest continuous tropical wetland: Integrative fire management is urgently needed for both biodiversity and humans. *J Environ Manage* 293: 112870. <https://doi.org/10.1016/j.jenvman.2021.112870>

- Grantham TE, Mezzatesta M, Newburn DA, Merenlender AM (2013) Evaluating tradeoffs between environmental flow protections and agricultural water security. *River Res Appl* 30: 315–328. <https://doi.org/10.1002/rra.2637>
- Guimarães EC, de Brito PS, Feitosa LM et al (2019) A new cryptic species of *Hyphessobrycon* Durbin, 1908 (Characiformes, Characidae) from the Eastern Amazon, revealed by integrative taxonomy. *Zoosyst Evol* 95(2): 345–360. <https://doi.org/10.3897/zse.95.34069>
- Hepp LU, Milesi SV, Biasi C, Restello RM (2010) Effects of agricultural and urban impacts on macroinvertebrates assemblages in streams (Rio Grande do Sul, Brazil). *Zoologia* 27: 106–113. <https://doi.org/10.1590/S1984-46702010000100016>
- Latini AO, Mormul RP, Giacomini HC et al (2021). Brazil's new fish farming Decree threatens freshwater conservation in South America. *Biol Conserv* 263: 109353. <https://doi.org/10.1016/j.biocon.2021.109353>
- Lewinsohn TM, Prado PI (2005) How many species are there in Brazil? *Conserv Biol* 19: 619–624. <https://doi.org/10.1111/j.1523-1739.2005.00680.x>
- Loiselle BA, Graham CH, Goerck JM, Ribeiro MC (2010) Assessing the impact of deforestation and climate change on the range size and environmental niche of bird species in the Atlantic forests, Brazil. *J. Biogeogr* 37(7): 1288–1301. <https://doi.org/10.1111/j.1365-2699.2010.02285.x>
- Maavara T, Chen Q, Van Meter K et al (2020) River dam impacts on biogeochemical cycling. *Nat Rev Earth Environ* 1(2): 103–116. <https://doi.org/10.1038/s43017-019-0019-0>
- Maffra MA, De Souza DC (2018) Barragens para irrigação: Aspectos jurídicos e ambientais da sua construção, operação e remoção. *Revista do CNMP* (7): 146–166.
- Malcolm JR, Liu C, Neilson RP, Hansen L, Hannah L (2006) Global warming and extinctions of endemic species from biodiversity hotspots. *Conserv Biol* 20(2): 538–548. <https://doi.org/10.1111/j.1523-1739.2006.00364.x>
- Marques OAV, Nogueira C, Martins M, Sawaya RJ (2010) Impactos potenciais das mudanças propostas no Código Florestal Brasileiro sobre os répteis brasileiros. *Biota Neotrop* 10: 39–41. <https://doi.org/10.1590/S1676-06032010000400004>
- Mittermeier RA, Câmara IG, Pádua MTJ, Blanck J (1990) Conservation in the Pantanal of Brazil. *Oryx* 24(2): 103–112. <https://doi.org/10.1017/S003060530003475X>
- MMA (2022). “Annex I. Official list of Brazilian fauna threatened with extinction”. Ordinance of Ministry of the Environment, Number 148, 7 July 2022. Retrieved in 14 August 2023.
- Nogueira DG, Valente-Neto F, Covich AP et al (2023) Riparian fruit input partially affects the distribution of stream fishes. *Hydrobiologia* 850: 3161–3172. <https://doi.org/10.1007/s10750-023-05242-2>
- Otoni FP, South J, Azevedo-Santos VM et al (2023) Editorial: Freshwater biodiversity crisis: Multidisciplinary approaches as tools for conservation. *Front Environ Sci* 11: 246. <https://doi.org/10.3389/fenvs.2023.1155608>
- Pelicice FM, Pompeu PS, Agostinho AA (2015) Large reservoirs as ecological barriers to downstream movements of Neotropical migratory fish. *Fish Fish* 16(4): 697–715. <https://doi.org/10.1111/faf.12089>
- Pelicice FM, Azevedo-Santos VM, Vitule JRS et al (2017) Neotropical freshwater fishes imperilled by unsustainable policies. *Fish Fish* 18(6): 1119–1133. <https://doi.org/10.1111/faf.12228>

- Pelicice FM, Agostinho AA, Akama A et al (2021) Large-scale degradation of the Tocantins-Araguaia River basin. *Environ Manage* 68: 445–452. <https://doi.org/10.1007/s00267-021-01513-7>
- Pelicice FM, Agostinho AA, Azevedo-Santos VM et al (2023) Ecosystem services generated by Neotropical freshwater fishes. *Hydrobiologia* 850: 2903–2926. <https://doi.org/10.1007/s10750-022-04986-7>
- Poff NL, Hart DD (2002) How dams vary and why it matters for the emerging science of dam removal. *BioScience* 52: 659–668. [https://doi.org/10.1641/0006-3568\(2002\)052\[0659:HDVAWI\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2002)052[0659:HDVAWI]2.0.CO;2)
- Pounds JA, Bustamante MR, Coloma MA et al (2006) Widespread amphibian extinctions from epidemic disease driven by global warming. *Nature* 439(7073): 161–167. <https://doi.org/10.1038/nature04246>
- Santana MLC, Carvalho FR, Teresa FB (2021) Broad and fine-scale threats on threatened Brazilian freshwater fish: variability across hydrographic regions and taxonomic groups. *Biota Neotrop* 21(2): e20200980. <https://doi.org/10.1590/1676-0611-BN-2020-0980>
- Simberloff D, Martin J-L, Genovesi P et al (2013) Impacts of biological invasions: what’s what and the way forward. *Trends Ecol Evol* 28: 58–66. <https://doi.org/10.1016/j.tree.2012.07.013>
- Souza AM, Marques DF, Döbereiner J, Dutra IS (2006). Esporos e toxinas de *Clostridium botulinum* dos tipos C e D em cacimbas no Vale do Araguaia, Goiás. *Pesq. Vet. Bras.* 26(3):133–138.
- Thomaz SM, Rosado A, Pires APF et al (2021) Nota técnica da Associação Brasileira de Limnologia para subsidiar as discussões sobre alterações das Áreas de Proteção Permanentes (APPs) urbanas propostas pelo PL 2.510/2019 (Câmara dos Deputados) e PL 1.869/2021 (Senado). ABLimno – Associação Brasileira de Limnologia. <https://doi.org/10.13140/RG.2.2.26832.17926>
- Vitule JRS, Umbria SC, Aranha JMR (2006) Introduction of the African catfish *Clarias gariepinus* (Burchell, 1822) into Southern Brazil. *Biol Invasions* 8: 677–681. <https://doi.org/10.1007/s10530-005-2535-8>
- Vitule JRS, Freire CA, Simberloff D (2009) Introduction of non-native freshwater fish can certainly be bad. *Fish Fish*, 10(1): 98–108. <https://doi.org/10.1111/j.1467-2979.2008.00312.x>
- Walters AW, Post DM (2011) How low can you go? Impacts of a low-flow disturbance on aquatic insect communities. *Ecol Appl* 21(1): 163–174. <https://doi.org/10.1890/09-2323.1>
- Wurtsbaugh WA, Paerl HW, Dodds WK (2019) Nutrients, eutrophication and harmful algal blooms along the freshwater to marine continuum. *Wiley Interdisciplinary Reviews: Water* 6(5): e1373. <https://doi.org/10.1002/wat2.1373>