

The text that follows is a PREPRINT.

O texto que segue é um PREPRINT.

Please cite as:

Favor citar como:

**Ziccardi, L.G., C.R. Thiersch, A.M. Yanai,
P.M. Fearnside & P.J. Ferreira-Filho.
2020. Forest-fire risk indices and zoning
of hazardous areas in Sorocaba, São
Paulo state, Brazil. *Journal of Forestry
Research* 31(2): 581-590.
<https://doi.org/10.1007/s11676-019-00889-x>**

[published online: 6 March 2019].

ISSN: 1007-662X (print version) ISSN: 1993-0607 (electronic version)

Copyright: Springer International Publishing AG. Part of Springer Nature

The original publication is available at:

O trabalho original está disponível em:

<http://www.springer.com/life+sciences/forestry/journal/11676>

<https://doi.org/10.1007/s11676-019-00889-x>

Journal of Forestry Research manuscript No. (will be inserted by the editor)
--

Forest-fire risk indices and zoning of hazardous areas in Sorocaba, São Paulo state, Brazil

Leonardo Guimarães Ziccardi ^{1,†}, Cláudio Roberto Thiersch ², Aurora Miho Yanai ¹, Philip Martin Fearnside ¹ and Pedro José Ferreira-Filho ²

Received: date / Accepted: date

Abstract Forest-fire risk indices and fire-risk maps are important tools used in protected areas to guide strategies for preventing and fighting fires. This study compares the performance of three fire-risk indices for accuracy in predicting fires in semi-deciduous forest fragments, creates a fire-risk map by integrating historical fire occurrences in a probabilistic density surface using the Kernel density estimator (KDE) in the municipality (county) of Sorocaba, São Paulo state, Brazil. The logarithmic Telicyn index, Monte Alegre formula (MAF) and enhanced Monte Alegre formula (MAF+) were employed using data for the period from 1 January 2005 to 31 December 2016. Meteorological data and numbers of fire occurrences were obtained from the National Institute of Meteorology (INMET) and the Institute for Space Research (INPE), respectively. Two performance measures were calculated: Heidke skill score (SS) and success rate (SR). The MAF+ index proved to be the most accurate for the study area, with values of skill score and success rate of 0.611 and 62.8%, respectively. The fire-risk map revealed two most susceptible areas with high (63 km²) and very high (47 km²) risk of fires in the municipality of Sorocaba. Identification of the best risk index and the generation of fire-risk maps can contribute to better planning and cost reduction in preventing and fighting forest fires.

Keywords Monte Alegre formula · risk map · forest protection · monitoring

¹ Department of Environmental Dynamics, National Institute for Research in Amazonia (INPA)
Av. André Araújo No. 2936, Manaus, Amazonas, CEP 69067-375 Brazil.

² Department of Environmental Sciences, Federal University of São Carlos (UFSCar)
Sorocaba, São Paulo, CEP 18052-780, Brazil.

[†] Tel.: +55-(11) 97574-5141; email: leonardo.g.ziccardi@gmail.com

1 Introduction

The forest fire regime is an important component in maintaining the function and structure of many terrestrial ecosystems, but it can also be considered a threat responsible for important economic and environmental impacts (e.g. economic losses in the forestry sector, degradation of land cover and changes in atmospheric composition) [13, 26]. If, on the one hand, forest fires can have a positive effect in terms of biodiversity and species richness [21, 29, 57], on the other hand can they impede hydrological processes and interfere both with the dynamic equilibrium of forests and with the carbon cycle [41, 15]. Considering the current scenario for global warming and expected increased frequency of occurrence of extreme events in the light of climate change, there is a tendency for future increase in the frequency of forest-fire outbreaks, mainly as a result of plant demographic processes that can change the growth of tree species and affect carbon sequestration by forests [9].

Monitoring meteorological parameters allows adoption of measures to reduce the potential damage of forest fires [17]. The main sources of ignition are human activities that are closely linked to the means of access to the forest, with increase in fire risk near roads in rural areas [12, 13]. Railways can also be sources of fire ignition, especially in the dry season. Sparks resulting from friction between the wheels of the train and the rails can start fires in dry vegetation, subsequently spreading to adjacent forests [27, 58].

An efficient plan to prevent and fight forest fires requires tools that include mapping areas that are most vulnerable to fire (i.e., fire-risk mapping) and the creation of forest-fire risk indices [30, 55]. A fire-risk map reveals the risk areas and facilitates the logistics for countermeasures by enabling rapid analysis of the situation for decision-making to prevent and combat forest fires [10]. Fire-risk indices show in advance the likelihood of forest-fire occurrence, and the interpretation of these values is linked to prevention plans and pre-suppression of fire [44, 55].

Several methods exist to interpolate historical ignition points and create a continuous wildfire risk map. The Kernel density estimation (KDE) is a nonparametric method that has been broadly used over the last two decades, especially after Koutsias et al. (2004), de la Riva et al. (2004) and Amatulli et al. (2007) explored how this technique is efficient to preserve a more realistic pattern of fire occurrence under broad pixel resolution [1, 22, 39], allowing appropriate forest fire-risk mapping [24].

Identifying areas of greatest fire risk makes it possible to adopt preventive measures in a timely fashion. These include construction of firebreaks, restriction of access to these locations in critical periods, reorganization of management and allocation of resources to strategic points [46, 50].

Fire-prevention plans in Brazil are mainly focused on protected areas. This is because these areas are fundamental to conserving biodiversity and maintaining ecological processes [16].

In addition to evaluating fire-risk indices, new models can be created to increase accuracy for the area under study. New tools can increase the efficiency of the resources allocated to protecting forests [55].

The objectives of the present study were to evaluate the accuracy of three fire-risk indices in semideciduous forest fragments, create a fire-risk map of the landscape in the municipality (county) of Sorocaba and determine the influence of roads and railways on the occurrence of fires.

2 Materials and Methods

2.1 Study site

This study was carried out in the rural portion of the municipality (county) of Sorocaba, São Paulo state, Brazil ($47^{\circ} 31' 50''$ W to $47^{\circ} 31'$ W and $23^{\circ} 34' 57''$ S to $23^{\circ} 35' 25''$ S)(Figure 1).




Fig. 1 Location of the municipality (county) of Sorocaba, São Paulo state, Brazil.

Sorocaba has 659,871 inhabitants and is one of the fastest growing cities in Brazil. Data released on July 1, 2017 as an estimate by the Brazilian Institute of Geography and Statistics (IBGE) show that, as compared to the estimate in 2016, the number of people living in Sorocaba increased by 1.13%, much more than the 0.77% growth in the country as a whole. In the last seven years, the city gained 73,246 new residents, and Sorocaba currently holds 13th place among Brazil's most populous municipalities (except for state capitals) [19].

The average elevation is 632 m with a maximum of 1028 m. The soil is classified as red latosol (Oxisol) and vegetation is that of a transition (ecotone) between Cerrado (central Brazilian savanna) and Atlantic forest, the former being very degraded and characterized by small fragments in secondary succession [14]. According to the Köppen classification, the climate is transitional Cwb (rainy and hot temperate with a moderately hot summer) to Cwa (seasonally dry with a warm to hot summer). Mean annual temperature is 22°C, and mean annual precipitation is 1310 mm [20].

2.2 Meteorological data and occurrence of fires

Daily records of average temperature (°C), relative humidity (%), wind speed (ms^{-1}) and precipitation (mm) used for the preparation of risk indices were obtained from the INMET (National Meteorology Institute) database for the period from 1 January 2005 to 31 December 2016 (Appendix A, Figure A1).

The hotspots that indicate fires, with their respective dates of occurrence and locations (latitude and longitude), were obtained from the INPE (National Institute of Space Research) database of MODIS satellite (AQUA_M-T) data with 1-km spatial resolution. Fire outbreaks were detected from the thermal signal composed of the wavelengths in the infrared range (Mid InfraRed-MIR) in a 500-m x 500-m pixel [40].

Occurrences were examined over the same period (1 January 2005 to 31 December 2016). Fire data were deleted in the case of duplicates, and 83 days were disregarded due to absence of weather data. The result was 4300 days of observation and 69 records of forest-fire occurrence in the study area.

2.3 Fire-risk map

As there are no uninhabited regions in the municipality, the risk map developed for the area was based on fire hazard associated with anthropogenic factors. This was done using a continuous density map of occurrences of outbreaks in a historical series of events [30, 31]. Annual outbreaks of fires were mapped based on their respective coordinates using ArcGIS software, UTM Zone 22 S projection and Datum SAD 69 (22s).

The Kernel density estimator (KDE) was used to determine critical areas for fire occurrence. This non-parametric density estimator produces a probabilistic density surface based on local information by superimposing a grid on

the data for each observed event [37, 18]. This tool adjusts for inaccuracies in the locatures of hotspots and is appropriate for manipulating spatial data at the scale of a municipality [22, 39].

2.4 Evaluation of the performance of fire-risk indices

The fire-risk indices are related to daily meteorological elements (Figure A2). They are considered to be in cumulative form, providing reliable results for the climatic characteristics of a region [49].

To determine the performance of each index it was necessary to define the limit separating presence and absence of fire risk, for which the rating scales of the indices were categorized in binary form. The indices indicate absence of risk when the calculated value is less than the lower limit of the medium risk class, while the presence of risk is indicated when the values are above this limit [36].

The Monte Alegre formula (MAF) is a cumulative index that uses two climatic variables for which values are easy to obtain, these being the number of days without rain (an indirect measure of precipitation) and the relative humidity at 13:00 h (1:00 pm). This index was developed based on data from the central portion of the state of Paraná [47], which is adjacent to the state of São Paulo where the current study was carried out. Use of MAF for determining the daily risk of forest fires is notable for being relatively easy [48].

$$MAF = \sum_{i=1}^n (100/H_i) \quad (1)$$

Where:

MAF	Monte Alegre formula
n	Number of consecutive days with precipitation less than 13 mm
i	Day number in a sequence of rainless days
H	Percentage of relative humidity measured at 13:00 h

The index is subject to modifications in the calculation according to the daily precipitation, which is necessary for obtaining cumulative values (Table B1). Estimates of the degree of danger associated with the calculated value of MAF should be interpreted according to a scale of risk (Table B2).

The enhanced Monte Alegre formula (MAF+) was developed from the inclusion of wind speed (ms^{-1}) as a variable modifying the MAF (Equation 2). As was the case for MAF, this formula was developed for central Paraná [35].

$$MAF+ = \sum_{i=1}^n (100/H_i) * e^{0.04*v} \quad (2)$$

Where:

MAF+	Enhanced Monte Alegre formula
n	Number of consecutive days with precipitation less than 13 mm
i	Day number in a sequence of rainless days
H	Percentage of relative humidity measured at 13:00 h
e	Base of natural logarithms (2.718282)
v	Wind speed (ms^{-1})

Since MAF+ is also cumulative in character, this index is subject to modifications in the calculation in accord with daily precipitation (Table B3). As is the case with MAF, the estimate of the degree of danger that is associated with the calculated value of MAF+ must be interpreted according to a scale of risk (Table B4). The variable corresponding to the wind speed (such as that at 13:00 h) is considered to be non-cumulative in the equation.

The logarithmic Telicyn Index (I) was developed in the former Union of Soviet Socialist Republics [51]. The two variables included in this index are air temperature and the dew point at 13:00 h. Like other indices considered in this study, the logarithmic Telicyn Index is also calculated cumulatively up to the moment of each precipitation event, after which a new calculation cycle begins:

$$I = \sum_{i=1}^n \log(T - r) \quad (3)$$

Where:

I	Telicyn index
T	Air temperature in °C
r	Dew point temperature in °C
log	Logarithm in base 10
n	Number of days without rain
i	Day number in a sequence of rainless days

The degree of danger related to the value calculated using Equation 3 must also be interpreted according to a scale of risk, and this index has only four classes of risk (Table B5). The logarithmic Telicyn index, the Monte Alegre formula (MAF) and the enhanced Monte Alegre formula (MAF+) were employed between 1 January 2005 and 31 December 2016. The predicted values for the occurrence of fires were obtained based on the scales for the three indices, the presence of risk being represented by the "medium," "high" and "very high" risk classes and the absence of risk by the "small" and "null" classes.

Comparison of the efficiency of the models was performed with two test parameters: the Heidke skill score (SS) and the success rate (SR). These parameters are based on a contingency table with observed and predicted values for occurrence of the events [34, 35, 36] (Table 1).

Table 1 Contingency table used to obtain the parameters that allow determination of ability to rate the skill score (SS) and success rate (SR). Adapted from [43].

Fires	Observed	Not observed	Total
Predicted	(a)	(b)	$N_2=a+b$
Not predicted	(c)	(d)	$N_4=c+d$
Total	$N_1=a+c$	$N_3=b+d$	$N=a+b+c+d$

Where:

N Total number of observations ($N = a + b + c + d$)

a Days on which occurrence of fires was predicted and they occurred (correct prediction)

b Days on which occurrence of fires was predicted and they did not occur (incorrect prediction)

c Days on which non-occurrence of fires was predicted and they occurred (incorrect prediction)

d Days on which non-occurrence of fires was predicted and they did not occur (correct prediction)

The following parameters were obtained from Table 1:

C Observed number of correct predictions (hits), where $C = a + d$

p Probability of having at least one event per day, where $p = N_1/N$

q Probability of exceeding the limit value of the index, in which $q = N_2/N$

E Expected number of hits, where $E = N * (1-p) * (1-q) + N * p * q$

SS Skill score, where $SS = (C-E)/(N-E)$

SR Success rate, where $SR = C/N$

3 Results

3.1 Outbreaks of forest fires

Of the 4300 days analyzed, 69 (1.6%) had occurrence of forest fires in the study area. The largest number of forest-fire outbreaks (15) occurred in 2014, while 2008 and 2013 had the fewest outbreaks (1) (Figure 2).




Fig. 2 Yearly distribution of outbreaks of forest fires from 2005 to 2016 in the municipality of Sorocaba, São Paulo state, Brazil.

Considering the monthly analysis of the cumulative total of outbreaks in the study period, August was the month with the highest number (23) of fire occurrences. The three-month period with the largest number of outbreaks was July, August and September (47) (Figure 3).




Fig. 3 Monthly distribution of outbreaks of forest fires from 2005 to 2016 in the municipality of Sorocaba, São Paulo state, Brazil. The red dashed lines represent the period with the greatest frequency of fires.

3.2 Performance of fire-risk indices

Performance of the indexes was quantified by the number of days from 2005 to 2016 for which fires were predicted and the number of days when fires were observed. Days in the study period were segregated into different classes in accord with each forest-fire risk index (Table 2).

Table 2 Distribution of days in fire-risk classes for each index from 2005 to 2016. Sorocaba, São Paulo, Brazil.

		Number of days per classes					
Risk		Absent		Present			
Index		Null	Small	Medium	High	Very high	Total
MAF	Numbers of days	413	727	1124	1127	909	4300
	Percentage	9,60%	16,90%	26,10%	26,20%	21,10%	
	Total (%)	26,50%		73,50%			100%
MAF +	Numbers of days	1140	1124	718	585	733	4300
	Percentage	26,50%	26,10%	16,70%	13,60%	17,10%	
	Total (%)	52,65%		47,35%			100%
Telicyn	Numbers of days	1570	433	341	1956	x	4300
	Percentage	36,50%	10,10%	7,90%	45,50%	x	
	Total (%)	46,60%		53,40%			x

The MAF showed the greatest numbers (3160 days or 73,50%) in the presence of fire-risk, considering the sum of "medium", "high" and "very high" classes. A tendency for a greater number of days in the classes for absence of risk was found for MAF+ (2264 days or 52.7%) as the sum of the "small" and "null" risk classes (Table 2).

The Telicyn risk index lacks the "very high" risk class and showed an intermediate number of observed days in classes that indicate the presence of risk, with 53.4%, considering the values of the "medium" and "high" risk classes. However, the percentage of days in low-risk classes was 46.6%, showing possible balance between classes (as was also the case for MAF+), probably depending on the climatic characteristics of the region.

Based on integration between the observed fire outbreaks and the adjusted scales of the risk indices, the values observed for fire occurrences were recorded in a contingency table. This was used to calculate skill score (SS) and success rate (SR) (Table 3).

Table 3 Number of days predicted by the indices for forest-fire occurrence and observed outbreaks in satellite data from 2005 to 2016 in the municipality of Sorocaba, São Paulo state, Brazil.

Risk index		Days observed	Days not observed	Total
MAF	Predicted	66	3063	3129
	Not predicted	3	1168	1171
	Total	69	4231	4300
MAF+	Predicted	53	1604	1657
	Not predicted	16	2627	2643
	Total	69	4231	4300
Telicyn	Predicted	64	2209	2273
	Not predicted	5	2022	2027
	Total	69	4231	4300

For calculation of SR, the days with hits or misses were quantified for each index. Hits were either days when occurrence of fire was predicted and one or

more outbreaks occurred or days when fire occurrence as not predicted and no outbreak occurred.

The SR result obtained by MAF+ (62.8%) was higher when compared to the other indices, with a SR value of 48.5% for the logarithmic Telicyn index and 28.7% for MAF (Table 4). The SS values (Table 4) mirror the results for SR, with MAF+ achieving the highest value (0.611), followed by Telicyn (0.468) and MAF (0.264).

Table 4 . Skill score (SS) and success rate (SR) values from 2005 to 2016 for fire-risk indices in the municipality of Sorocaba, São Paulo state, Brazil.

Risk index	SS	SR
MAF	0.264	28.70%
MAF+	0.611	62.80%
Telicyn	0.468	48.50%

3.3 Fire-risk map

The risk map drawn from the smoothed density of fire outbreaks in the study period (2005 to 2016) identifies areas favorable for forest fires (Figure 4). The values generated by the algorithms for calculating Kernel density were categorized into five classes for classifying risk levels. Interpretation of risk levels (very low, low, moderate, high and very high) is a key step in delimitating areas that are most vulnerable to fire as part of a prevention plan [13].




Fig. 4 Fire-risk map for the municipality of Sorocaba, São Paulo state, Brazil.

4 Discussion

Many forest-fire risk indices derived from daily meteorological variables have been created around the world based on the assumption that climate-related variables are the most important in determining fire risk [56]. The most frequently cited fire-risk indices in the literature are the Russian Nesterov Index [32], the Russian Logarithmic Index of Telicyn [51], the U.S. National Fire Danger Rating System [11], the Swedish Angstrom index [7], the Canadian Forest Fire Danger Rating System (CFFDRS) [52] and the Argentine index of Rodriguez and Moretti [42].

In Brazil, FMA [47] is the index that is most used by forestry and environment protection companies [54]. However, some studies have revealed that this is not the most suitable index for certain Brazilian regions, especially the north [45], northeast [54], mid-west [31] and even the southeast region [25], which is close to where the index was created. This is because most of the

forest fire hazard indices based on empirical models are suitable only for local application because of the specific vegetation and climate from where they were developed [53].

The SS value obtained for MAF+ in the present study (0.611) was higher than the values found by Nunes [33] for the same index in Paraná for the period from 1998 to 2003 in the municipalities of Cambará, Telêmaco Borba, Guarapuava, Pinhais, Campo Mourão, Cascavel and Londrina (0.088, 0.117, 0.133, 0.283, 0.302, 0.334 and 0.338, respectively). The SR values we found for MAF (28.7%) and Telicyn (48.5%) in Sorocaba were below that for Guarapuava, which had the smallest value (49.3%) of the 13 municipalities evaluated by Nunes [33] in the state of Paraná.

Much of the rural area of Sorocaba is under eucalyptus plantations, and the area of these plantations rose from 233,406 ha in 2002 [23] to 323,478 ha in 2008 [5], an increase of 38.6% over this relatively short period. These areas have high risk of fires because, in addition to the availability of wood, a blanket of combustible material is offered by a continuous deposition of leaves and twigs on the soil surface [4]. Generally, the boundaries between zones of high and low risk of fires are consistent with the limits between continuous and fragmented forest [38].

Compared to the other indices, MAF+ showed the best performance in tests of SS and SR, as has also been reported for the period from 2003 to 2006 in predicting outbreaks of fire in areas with eucalyptus cultivation in the northern portion of the state of Espírito Santo, Brazil, with values of 0.18 for SR and 53.5%, for SS [4].

MAF+ includes wind speed as a variable, which differs from the original formula (MAF). With inclusion of this variable, MAF+ reflects, in addition to the probability of ignition, the potential for fires to spread [35].

The forest-fire risk map revealed two areas with the highest occurrences of outbreaks, the northwest and southeast portions of the study area. Of the entire area of the municipality of Sorocaba (343.1 km²), 13.7% was classified as "very high risk" (47 km²) and 18.5% was classified as "high risk" (63 km²).

The northwestern portion of the municipality has eucalyptus plantations and lies next to the Ipanema National Forest; the northwest portion of the municipality also has a large number of roads. The high productivity of commercial plantations in these areas leads to a high concentration of biomass, which raises the risk of forest fires [6].

In the southeastern portion of the municipality, the large number of outbreaks recorded can be linked, in part, to the stretch of railway in association with a green area. Presence of the railway may therefore facilitate ignition and increase the risk of fires in adjacent vegetation in association with wind patterns [8].

The spatial statistical analysis of weights of evidence is able to capture the effect of distances of roads and railroads on fire outbreaks. It is a probabilistic method based on a Bayesian approach in log-linear form and is applicable when sufficient data are available to estimate the relative influence of different scenarios for the factors considered in the analysis [3]. However, the low spatial

resolution of MODIS does not allow discussion of distances of less than 1 km, making this analysis unfeasible for this data set.

One suggestion for future studies aiming to use this analysis is to obtain the hotspots that indicate fire through the LANDSAT 8 thermal band. The spatial resolution of this reference satellite is 30 m (i.e., with much more spatial detail than MODIS), but the temporal resolution is 16 days, leading to the omission of many fire outbreaks.

The scenario for fire in a given location is linked to the socio-economic, political and environmental context of the region. There is a strong connection between the fire regime and territorial dynamics at different temporal scales [28]. Recent studies also point to the existence of a relationship between the type of vegetation and the frequency and intensity of forest fires in any given location [2].

Because fire-prevention planning requires monitoring of where and when a fire is likely to occur [24], the forest-fire risk map and fire-risk indices are two objective tools that should be used together for efficient pre-fire planning. While the fire-risk map helps managers in planning fire prevention strategies for showing the most fire-susceptible areas (based on historical fire occurrences), the indices indicate which days are most likely to the occurrence of forest fires (based on daily meteorological factors). Determining the best index for application in municipalities that do not have an index specific to their location is fundamental for planning forest-fire prevention.

MAF+ had the best result among the indices examined in this study. However, Brazil is a country with a wide diversity of climates, which is reflected in the different results of the various indices of forest-fire risk in different regions of the country.

5 Conclusions

Depending on climatic patterns in the region under study, one risk index may be more suitable than another for the prediction of fire events. The MAF+ index had the best performance in the municipality of Sorocaba, with a skill score (SS) value of 0.611 and a success rate (SR) of 62.3%. This indicates that wind speed is an important variable and should be considered for determining the degree of fire hazard in this region.

By developing a risk map, it was possible to identify the areas that are favorable for occurrence of forest fires. These areas are mainly in the north-western and southeastern portions of the municipality. The risk of forest fires in Sorocaba is associated with local climatic conditions, with the network of roads and with the railway that crosses the municipality.

Acknowledgements: We thank the National Institute for Research in Amazonia (INPA: PRJ15.125), the National Institute of Science and Technology for the Environmental Services of Amazonia (INCT-SERVAMB), the National Council for Scientific and Technological Development (CNPq/PCI

Program Proc. 304130/2013-3 and 301183/2015-5; CNPq: Proc. 304020/2010-9; 573810/2008-7) and the Foundation for Support of Research in Amazonas State (FAPEAM: Proc. 708565) for financial and logistical support. We owe personal thanks to Alberto Vicentini, Nathan Goncalves, Ricardo Perdizes and Vitor Alves.

This article is a contribution of the Brazilian Research Network on Global Climate Change, FINEP/ Rede CLIMA Grant Number 01.13.0353-00.

The authors declare no conflict of interest. The funding sponsors had no role in the design of the study, in the collection, analyses do, or interpretation of data, in the writing of the manuscript, and in the decision to publish the results.

References

1. Amatulli G, Pérez-Cabello F, de la Riva J (2007) Mapping lightning/human-caused wildfires occurrence under ignition point location uncertainty. *Ecological Modelling* 200(3):321–333
2. Bernier PY, Gauthier S, Jean PO, Manka F, Boulanger Y, Beaudoin A, Guindon L (2016) Mapping local effects of forest properties on fire risk across Canada. *Forests* 7(8):157
3. Bonham-Carter GF (2014) Geographic information systems for geoscientists: modelling with GIS, vol 13. Elsevier
4. Borges TS, Fiedler NC, dos Santos AR, Loureiro EB, Mafia RG (2011) Desempenho de alguns índices de risco de incêndios em plantios de eucalipto no norte do Espírito Santo [performance of some fire risk indices in eucalyptus plantations in northern Espírito Santo]. *Floresta e Ambiente* 18(2):153 – 159
5. Canavesi V, Célia R, Alvalá R, Cunha AP, Cunha A (2011) Análise espaço-temporal dos plantios de Eucalyptus spp. no Estado de São Paulo [spatiotemporal analysis of Eucalyptus spp. plantations in the state of São Paulo]. *Anais XV Simpósio Brasileiro de Sensoriamento Remoto - SBSR*, Curitiba, PR, Brazil, INPE p.2113
6. Castro F, Tudela A, Sebastià MT (2003) Modeling moisture content in shrubs to predict fire risk in Catalonia (Spain). *Agricultural and Forest Meteorology* 116(1):49–59
7. Chandler C, Cheney P, Thomas P, Traubaud L, Williams D, et al (1983) Fire in forestry. Volume 1. Forest fire behavior and effects. Volume 2. Forest fire management and organization. John Wiley & Sons, Inc.
8. Chuvieco E, Aguado I, Yebra M, Nieto H, Salas J, Martín MP, Vilar L, Martínez J, Martín S, Ibarra P, et al (2010) Development of a framework for fire risk assessment using remote sensing and geographic information system technologies. *Ecological Modelling* 221(1):46–58
9. Collins L, Penman T, Ximenes FA, Binns D, York A, Bradstock R (2014) Impacts of frequent burning on live tree carbon biomass and demography in post-harvest regrowth forest. *Forests* 5(4):802–821

10. Dalcumune MAB, Santos A (2005) Mapeamento de índice de risco de incêndio para a região da grande Vitória/ES, utilizando imagens do satélite Landsat para o ano de 2002 [mapping of the fire risk index for the region of greater Vitória/ES using images from the Landsat satellite for the year 2002]. *Anais do XII Simpósio Brasileiro de Sensoriamento Remoto* pp 1485–1492
11. Deeming JE, Burgan RE, Cohen JD (1977) The national fire-danger rating system, 1978. Tech. rep.
12. Díaz-Delgado R, Lloret F, Pons X (2004) Spatial patterns of fire occurrence in Catalonia, NE, Spain. *Landscape Ecology* 19(7):731–745
13. Duarte L, Teodoro AC (2016) An easy, accurate and efficient procedure to create forest fire risk maps using the SEXTANTE plugin Modeler. *Journal of Forestry Research* 27(6):1361–1372
14. EMBRAPA (2006) Empresa brasileira de pesquisa agropecuária [brazilian agricultural research corporation]. <http://www.embrapa.br>, Accessed: 9 March 2015.
15. Fearnside PM, Barbosa RI (1999) Incêndios na Amazônia brasileira: estimativa da emissão de gases do efeito estufa pela queima de diferentes ecossistemas de Roraima na passagem do evento El Niño (1997/98) [fires in the Brazilian Amazon: estimation of the emission of greenhouse gases by the burning of different ecosystems of Roraima in the passage of the El Niño (1997/98)]. *Acta Amazonica* 29(4):513–534
16. Fiedler NC (2004) Incêndios florestais no Parque Nacional da Serra da Canastra: desafios para a conservação da biodiversidade [forest fires in Serra da Canastra National Park: challenges for biodiversity conservation]. *Ciência Florestal* 14(2):157–168
17. França Tetto A, Batista AC, Soares Nunes JR, Viana Soares R (2010) Subsídios à prevenção e combate a incêndios florestais com base no comportamento da precipitação pluviométrica na Floresta Nacional de Irati, Paraná [subsidies for the prevention and combat of forest fires based on rainfall behavior in the Irati National Forest, Paraná]. *Ciência Florestal* 20(1):33–43
18. Gatrell AC, Bailey TC, Diggle PJ, Rowlingson BS (1996) Spatial point pattern analysis and its application in geographical epidemiology. *Transactions of the Institute of British Geographers*; 21:256–274
19. IBGE (2017) Instituto Brasileiro de Geografia e Estatística [Brazilian Institute of Geography and Statistics]. <http://www.ibge.gov.br>, Accessed: 13 December, 2017.
20. Ikematsu P, da Silva AM, de Paula FP, Nogueira DP, Silveira FM, Alves SH, Bomback M (2007) Dimensionamento e estudo dos fatores condicionantes de duas voçorocas localizadas no município de Sorocaba (SP) [dimensioning and study of the conditioning factors of two gullies located in the municipality of Sorocaba (SP)]. *Caminhos de Geografia* 8(24):76–85.
21. Keeley JE, Lubin D, Fotheringham C (2003) Fire and grazing impacts on plant diversity and alien plant invasions in the southern Sierra Nevada. *Ecological Applications* 13(5):1355–1374

22. Koutsias N, Kalabokidis KD, Allgöwer B (2004) Fire occurrence patterns at landscape level: beyond positional accuracy of ignition points with kernel density estimation methods. *Natural Resource Modeling* 17(4):359–375
23. Kronka FJ, Nalon MA, Matsukuma CK, et al (2002) Inventário florestal das áreas reflorestadas no Estado de São Paulo [forest inventory of reforested areas in the state of São Paulo]. SMA/IF, p 184
24. Kuter N, Yenilmez F, Kuter S (2011) Forest fire risk mapping by kernel density estimation. *Croatian Journal of Forest Engineering* 32(2):599–610
25. Mafalda V, Torres F, Ribeiro G (2009) Eficiência de índices de perigo de incêndios baseados em elementos climáticos no município de Juiz de Fora - MG [efficiency of fire hazard indexes based on climatic elements in the municipality of Juiz de Fora - MG]. XIII SBGFA-Simpósio Brasileiro de Geografia Física Aplicada, Viçosa, MG
26. Malowerschnig B, Sass O (2014) Long-term vegetation development on a wildfire slope in Innerzwain (Styria, Austria). *Journal of Forestry Research* 25(1):103–111
27. Martell DL, Otukol S, Stocks BJ (1987) A logistic model for predicting daily people-caused forest fire occurrence in Ontario. *Canadian Journal of Forest Research* 17(5):394–401
28. Montiel Molina C, Galiana-Martín L (2016) Fire scenarios in Spain: A territorial approach to proactive fire management in the context of global change. *Forests* 7(11):273
29. Moretti M, Duelli P, Obrist MK (2006) Biodiversity and resilience of arthropod communities after fire disturbance in temperate forests. *Oecologia* 149(2):312–327
30. Morgan P, Hardy CC, Swetnam TW, Rollins MG, Long DG (2001) Mapping fire regimes across time and space: understanding coarse and fine-scale fire patterns. *International Journal of Wildland Fire* 10(4):329–342
31. Narciso M, Soriano B, Daniel O, Máximo F (2011) Proposta de método para seleção de indicador de risco de incêndio por região [proposal of method for selection of fire risk indicator by region]. Embrapa Pantanal-Boletim de Pesquisa e Desenvolvimento (INFOTECA-E)
32. Nesterov V (1949) Fire frequency index and method of its estimation. Goslesbumaga, Moscow
33. Nunes JRS (2007) Sistema integrado de controle de incêndios florestais para o estado do paraná [integrated forest fire control system for the state of paraná]. PhD thesis, UFPR
34. Nunes JRS (2008) Desempenho da Fórmula de Monte Alegre (FMA) e da Fórmula de Monte Alegre alterada (FMA+) no Distrito Florestal de Monte Alegre, município de Telêmaco Borba, Paraná [performance of the Monte Alegre Formula (MAF) and Monte Alegre modified formula (FMA+) in the Monte Alegre Forest District, Telêmaco Borba, Paraná], pp. 70. PhD thesis, UFPR
35. Nunes JRS, Soares RV, Batista AC (2006) FMA+ - um novo índice de perigo de incêndios florestais para o Estado do Paraná, Brasil [FMA+ - a new forest fire hazard index for the State of Paraná, Brazil]. *Floresta* pp.

- 36(1)
36. Nunes JRS, Fier ISN, Soares RV, Batista AC (2010) Desempenho da fórmula de Monte Alegre (FMA) e da fórmula de Monte Alegre alterada (FMA+) no Distrito Florestal de Monte Alegre [performance of the Monte Alegre formula (FMA) and Monte Alegre modified formula (FMA+) in the Forest District of Monte Alegre]. *Revista Floresta* 40(2):319 – 326
 37. Parzen E (1962) On estimation of a probability density function and mode. *The Annals of Mathematical Statistics* 33(3):1065–1076
 38. Portier J, Gauthier S, Leduc A, Arseneault D, Bergeron Y (2016) Fire regime along latitudinal gradients of continuous to discontinuous coniferous boreal forests in eastern Canada. *Forests* 7(10):211
 39. De la Riva J, Pérez-Cabello F, Lana-Renault N, Koutsias N (2004) Mapping wildfire occurrence at regional scale. *Remote Sensing of Environment* 92(3):363–369
 40. Roberts GJ, Wooster MJ (2008) Fire detection and fire characterization over Africa using Meteosat SEVIRI. *IEEE Transactions on Geoscience and Remote Sensing* 46(4):1200–1218
 41. Robinne FN, Miller C, Parisien MA, Emelko MB, Bladon KD, Silins U, Flannigan M (2016) A global index for mapping the exposure of water resources to wildfire. *Forests* 7(1):22
 42. Rodriguez N, Moretti A (1988) Índice de peligro de propagación de incendios forestales [index of propagation danger of forest fires]. In: 6. Congreso Forestal Argentino. Santiago del Estero (Argentina). 16-20 Ago 1988., vol 3, pp 704–709
 43. Sampaio OB (1999) Análise da eficiência de quatro índices na previsão de incêndios florestais para a região de Agudos-SP [analysis of the efficiency of four indices in the prediction of forest fires for the region of Agudos-SP]. PhD thesis, UFPR
 44. SantAnna CM, Fiedler NC, Minette LJ (2007) Controle de incêndios florestais [control of forest fires]. *Alegre: Suprema* p 152
 45. da Silva ID, Pontes Jr AC (2011) Elaboração de um fator de risco de incêndios florestais utilizando lógica fuzzy [development of a risk factor for forest fires using fuzzy logic]. *Biomatematica* 21:113–128
 46. Smith DT (1998) Mapping the bushfire danger of Hobart. PhD thesis, University of Tasmania
 47. Soares R (1972) Determinação de um índice de perigo de incêndio para a região centro paranaense, Brasil [determination of a fire hazard index for the central region of Paraná, Brazil]. *turrialba, costa rica*, 72 pp. Master's thesis, CATIE/IICA
 48. Soares RV (1998) Desempenho da "Fórmula de Monte Alegre" índice brasileiro de perigo de incêndios florestais [performance of the "Monte Alegre Formula" Brazilian index of forest fire hazard]. *Cerne* 4(1):87 – 89
 49. Soares RV, Batista AC (2007) Incêndios florestais: controle, efeitos e uso do fogo [Forest fires: control, effects and use of fire]. 223-238, Universidade Federal do Paraná

-
50. Sousa C (2000) Relatório do projeto de cartografia de risco de incêndio florestal-CRIF 2^a fase [report of the forest fire risk cartography project-CRIF 2nd phase]. Available at: www.terraviva.pt (Accessed: april 19, 2017) 17
51. Telicyn G (1970) Logarithmic index of fire weather danger for forests. *Lesnoe Khozyaistvo* (11):58–9
52. Van Wagner C, Forest P, et al (1987) Development and structure of the Canadian forest fireweather index system. In: *Can. For. Serv., Forestry Tech. Rep.*, Citeseer
53. Wastl C, Schunk C, Leuchner M, Pezzatti GB, Menzel A (2012) Recent climate change: long-term trends in meteorological forest fire danger in the Alps. *Agricultural and Forest Meteorology* 162:1–13
54. White BLA, White LAS, Ribeiro GT, Fernandes PAM (2013) Development of a fire danger index for eucalypt plantations in the northern coast of Bahia, Brazil. *Floresta* 43(4):601–610
55. White LAS, White BLA, Ribeiro GT (2015) Evaluation of forest fire danger indexes for eucalypt plantations in Bahia, Brazil. *International Journal of Forestry Research* 2015(613736):6
56. Whitlock C, Shafer SL, Marlon J (2003) The role of climate and vegetation change in shaping past and future fire regimes in the northwestern us and the implications for ecosystem management. *Forest Ecology and Management* 178(1):5–21
57. Wohlgemuth T, Moretti M, Conedera M, Moser B (2006) Ecological resilience after fire in mountain forests of the central Alps. *Forest Ecology and Management* 234(1):S200
58. Wotton BM, Nock CA, Flannigan MD (2010) Forest fire occurrence and climate change in Canada. *International Journal of Wildland Fire* 19(3):253–271

A Appendix

A.1




Fig. 5 Relationships between environmental variables, where $R^2_{ad.}$ is the adjusted R-squared. Data are from Brazil's National Institute of Meteorology (INMET).

548 A.2




Fig. 6 Relationships between environmental variables and fire indices, where $R^2_{ad.}$ is the adjusted R-squared.

B Appendix

B.1

Table 5 Modification of calculation of the MAF on the basis of daily precipitation.

Daily precipitation (mm)	Change in calculation
≤ 2.4	None
2.5 to 4.9	30% reduction in the MAF calculated for the previous day and sum of (100/H) for the day
5.0 to 9.9	60% reduction of the MAF calculated for the previous day and sum of (100/H) for the day
10.0 to 12.9	80% reduction of the MAF calculated for the previous and sum of (100/H) for the day
> 12.9	Calculation interrupted (MAF = 0) and summing resumed the following day

B.2

Table 6 Scale of risk classes associated with the calculated values of MAF.

Calculated value of MAF	Forest-fire risk
≤ 1.0	Null
1.1 to 3.0	Small
3.1 to 8.0	Medium
8.1 to 20.0	High
> 20.0	Very high

B.3

Table 7 Modification of calculation of MAF+ as a function of daily precipitation.

Daily precipitation (mm)	Change in calculation
≤ 2.4	None
2.5 to 4.9	30% reduction in the MAF calculated for the previous day and sum of (100/H) for the day
5.0 to 9.9	60% reduction of the MAF calculated for the previous day and sum of (100/H) for the day
10.0 to 12.9	80% reduction of the MAF calculated for the previous and sum of (100/H) for the day
> 12.9	Calculation interrupted (MAF = 0) and summing resumed the following day

553 B.4

Table 8 Scale of risk classes associated with the calculated values of MAF+.

Calculated value of MAF	Forest-fire risk
≤ 3.0	Null
3.1 to 8.0	Small
8.1 to 14.0	Medium
14.1 to 24.0	High
> 24.0	Very high

554 B.5

Table 9 Scale of risk classes associated with the calculated values of Telicyn.

Calculated value of Telicyn	Forest-fire risk
≤ 2.0	Null
2.1 to 3.5	Small
3.6 to 5.0	Medium
> 5.0	High